

On invariant CCC σ –ideals.

Jan Kraszewski

Mathematical Institute,

University of Wrocław,

pl. Grunwaldzki 2/4,

50-384 Wrocław, Poland

(e-mail: kraszew@math.uni.wroc.pl)

Abstract

We re-read Reclaw's proof from [6] on invariant CCC σ –ideals of subsets of reals and obtain a reasonably stronger corollary for such ideals on the Cantor space.

1. Preliminaries. In 1998 Reclaw in [6] investigated cardinal invariants of CCC σ –ideals of subsets of reals. In particular, he showed that if such a σ –ideal \mathcal{J} is invariant, then $\mathfrak{p} \leq \text{non}(\mathcal{J})$, where \mathfrak{p} is a pseudointersection number (cf. [8] for more details). In this paper we analyze his proof and get an apparently stronger result for σ –ideals of subsets of the Cantor space 2^ω .

We use standard set-theoretical notation and terminology derived from [1]. Let us remind that the cardinality of the set of all real numbers is denoted by \mathfrak{c} . The cardinality of a set X is denoted by $|X|$. By $[\omega]^\omega$ we denote the family of all infinite subsets of ω . If $\varphi : X \rightarrow Y$ is a function then $\text{rng}(\varphi)$ denotes the range of φ .

Let $(G, +)$ be an abelian Polish (i.e. separable, completely metrizable, without isolated points) group and let \mathcal{J} be a σ –ideal of subsets of G (we assume from now on that \mathcal{J} is proper and contains all singletons). We will consider that \mathcal{J} is invariant, that is for every $A \subseteq G$ and $g \in G$ we have $A + g = \{a + g : a \in A\} \in \mathcal{J}$ and $-A = \{-a : a \in A\} \in \mathcal{J}$. Moreover, we will assume that the σ –ideal \mathcal{J} has a Borel basis i.e. every set from \mathcal{J} is contained in a certain Borel set from the ideal.

We say that \mathcal{J} is CCC (countable chain condition) if the quotient Boolean algebra $\mathcal{B}(G)/\mathcal{J}$ is CCC, where $\mathcal{B}(G)$ is the σ –algebra of all Borel subsets of G .

We define the following cardinal invariants of \mathcal{J} .

$$\begin{aligned} \text{non}(\mathcal{J}) &= \min\{|B| : B \subseteq G \wedge B \notin \mathcal{J}\}, \\ \text{cov}_t(\mathcal{J}) &= \min\{|T| : T \subseteq G \wedge (\exists A \in \mathcal{J}) A + T = G\}, \end{aligned}$$

2000 *Mathematics Subject Classification*: 03E05, 03E17.

Key words and phrases: invariant σ –ideal, CCC, cardinal invariant.

We define also an operation on the σ -ideal \mathcal{J} (it was introduced by Seredyński in [7], who denoted it by \mathcal{J}^*)

$$s(\mathcal{J}) = \{A \subseteq G : (\forall B \in \mathcal{J})(\exists g \in G)(A + g) \cap B = \emptyset\}.$$

If we apply these operations to the σ -ideals of meagre sets \mathcal{M} and of null sets \mathcal{N} we obtain strongly null sets $s(\mathcal{M})$ and strongly meager sets $s(\mathcal{N})$. The following is well-known

$$\text{non}(s(\mathcal{J})) = \text{cov}_t(\mathcal{J}).$$

We define

$$Pif = \{f : f \text{ is a function} \wedge \text{dom}(f) \in [\omega]^\omega \wedge \text{rng}(f) \subseteq 2\}.$$

If $f \in Pif$ then we put

$$[f] = \{x \in 2^\omega : f \subseteq x\}.$$

Let \mathbb{S}_2 denotes the σ -ideal of subsets of the Cantor space 2^ω , which is generated by the family $\{[f] : f \in Pif\}$. It was thoroughly investigated in [2] and [4]. We recall some properties of \mathbb{S}_2 , which were proved in [2].

Fact 1.1 \mathbb{S}_2 is a proper, invariant σ -ideal which contains all singletons and has a Borel basis. Every $A \in \mathbb{S}_2$ is both meager and null. Moreover, there exists a family of size \mathfrak{c} of pairwise disjoint Borel subsets of 2^ω that do not belong to \mathbb{S}_2 . Hence \mathbb{S}_2 is not CCC. \square

Let A, S be two infinite subsets of ω . We say that S splits A if $|A \cap S| = |A \setminus S| = \omega$. Let us recall a cardinal number related with a notion of splitting, introduced by Malychin in [5], namely

$$\aleph_{0-\mathfrak{s}} = \min\{|\mathcal{S}| : \mathcal{S} \subseteq [\omega]^\omega \wedge (\forall \mathcal{A} \in [[\omega]^\omega]^\omega)(\exists S \in \mathcal{S})(\forall A \in \mathcal{A}) S \text{ splits } A\}.$$

More about cardinal numbers connected with the relation of splitting can be found in [3].

2. Reclaw's proof revisited. In [6] Reclaw proved a theorem, which can be generalized as follows.

Theorem 2.1 Let \mathcal{I} and \mathcal{J} be two σ -ideals of subsets of an abelian Polish group G , which are invariant and have Borel bases. If \mathcal{I} is CCC then

$$\mathcal{J} \cap s(\mathcal{J}) \subseteq \mathcal{I}.$$

Proof. (Reclaw) Let $X \in \mathcal{J} \cap s(\mathcal{J})$. Assume that $X \notin \mathcal{I}$. We construct a sequence $\{F_\alpha : \alpha < \omega_1\}$ of Borel sets from \mathcal{J} and a sequence $\{t_\alpha : \alpha < \omega_1\}$ of elements of G . Let $t_0 = 0$ and F_0 be any Borel set from \mathcal{J} containing X . Suppose that we have constructed F_β and t_β for $\beta < \alpha$. Then from the definition of $s(\mathcal{J})$ there exists $t_\alpha \in G$ such that

$$(X + t_\alpha) \cap \bigcup_{\beta < \alpha} F_\beta = \emptyset.$$

As F_α we take any Borel set from \mathcal{J} containing $\bigcup_{\beta < \alpha} F_\beta \cup (X + t_\alpha)$.

Let $G_\alpha = F_\alpha \setminus \bigcup_{\beta < \alpha} F_\beta$. Thus $\{G_\alpha : \alpha < \omega_1\}$ is a family of pairwise disjoint Borel sets such that none of them belongs to \mathcal{I} , as $G_\alpha \supseteq X + t_\alpha$ and \mathcal{I} is invariant. Hence \mathcal{I} is not CCC, a contradiction. \square

Corollary 2.2 *Let \mathcal{I} and \mathcal{J} be as above. If \mathcal{I} is CCC then*

$$\min\{\text{non}(\mathcal{J}), \text{cov}_t(\mathcal{J})\} \leq \text{non}(\mathcal{I}).$$

Proof. It is enough to observe that $\mathcal{J} \subseteq \mathcal{I}$ implies $\text{non}(\mathcal{J}) \leq \text{non}(\mathcal{I})$. \square

Corollary 2.3 *Let \mathcal{I} be a σ -ideal of subsets of the Cantor space 2^ω (endowed with a standard group structure), which is invariant and has a Borel basis. If \mathcal{I} is CCC then*

$$\aleph_{0-\mathfrak{s}} \leq \text{non}(\mathcal{I}).$$

Proof. In [2] it was proved that $\text{non}(\mathbb{S}_2) = \aleph_{0-\mathfrak{s}}$ and in [4] it was proved that $\text{cov}_t(\mathbb{S}_2) = \mathfrak{c}$. So it is enough to apply Corollary 2.2 for $G = 2^\omega$ and $\mathcal{J} = \mathbb{S}_2$. \square

Question. Let \mathcal{I} be an invariant CCC σ -ideal of subsets of the real line \mathbb{R} . Is the inequality $\aleph_{0-\mathfrak{s}} \leq \text{non}(\mathcal{I})$ still true?

References

- [1] T. Bartoszyński, H. Judah, *Set Theory: On the structure of the real line*, A. K. Peters, Wellesley, Massachusetts 1995.
- [2] J. Cichoń, J. Kraszewski, *On some new ideals on the Cantor and Baire spaces*, Proc. Am. Math. Soc., **126** (1998) 1549–1555.
- [3] A. Kamburelis, B. Węglorz, *Splittings*, Arch. Math. Logic, **35** (1996) 263–277.
- [4] J. Kraszewski, *Transitive properties of the ideal \mathbb{S}_2* , Real Anal. Ex., 29(2) (2003/2004), 629–638.
- [5] V.I. Malychin, *Topological properties of Cohen generic extension*, Trans. Mosc. Math. Soc., **52** (1990) 1–32.
- [6] I. Reclaw, *On cardinal invariants for CCC σ -ideals*, Proc. Am. Math. Soc., **126** (1998) 1173–1175.
- [7] W. Sieredyński, *Some operations related with translation*, Colloq. Math., **57** (1989) 203–219.
- [8] J. E. Vaughan, *Small uncountable cardinals and Topology*, in: *Open Problems in Topology*, North-Holland, Amsterdam etc., (1990), 195–218