

## A CHAOTIC DECOMPOSITION FOR GENERALIZED STOCHASTIC PROCESSES WITH INDEPENDENT VALUES

BY

SUMAN DAS (SWANSEA) AND EUGENE LYTVYNOV\* (SWANSEA)

*Abstract.* We extend the result of Nualart and Schoutens on chaotic decomposition of the  $L^2$ -space of a Lévy process to the case of a generalized stochastic processes with independent values.

**2000 AMS Mathematics Subject Classification:** Primary: 60H05, 60H40; Secondary: 60B05.

**Key words and phrases:** Chaotic decomposition, generalized stochastic process, Lévy process.

### 1. INTRODUCTION

Among all stochastic processes with independent increments, essentially only Brownian motion and Poisson process have a chaotic representation property. The latter property means that, by using multiple stochastic integrals with respect to the centered stochastic process, one can construct a unitary isomorphism between the  $L^2$ -space of the process and a symmetric Fock space. In the case of a Lévy process, several approaches have been proposed in order to construct a Fock space-type realization of the corresponding  $L^2$ -space. In this paper, we will be concerned with the approach of Nualart and Schoutens [9], who constructed a representation of every square integrable functional of a Lévy process in terms of orthogonalized Teugels martingales. Recall that, for a given Lévy process  $(X_t)_{t \geq 0}$ , its  $k$ -th order Teugels martingale is defined by centering the power jump process

$$X_t^{(k)} := \sum_{0 < s \leq t} (\Delta X_s)^k, \quad k \in \mathbb{N}.$$

For numerous applications of this result, see e.g. [6] and [10]. We also refer to [7] for an extension of this result to the case of a Lévy process taking values in  $\mathbb{R}^d$ , and

---

\* E.L. acknowledges the financial support of the Polish National Science Centre, grant no. 2012/05/B/ST1/00626, and of the SFB 701 “Spectral structures and topological methods in mathematics”, Bielefeld University.

to [1] and [3] for a Nualart–Schoutens-type decomposition for noncommutative (in particular, free) Lévy processes.

The aim of this note is to extend the Nualart–Schoutens decomposition to the case of a generalized stochastic process with independent values. Consider a standard triple  $\mathcal{D} \subset L^2(\mathbb{R}^d, dx) \subset \mathcal{D}'$ , where  $\mathcal{D} = C_0^\infty(\mathbb{R}^d)$  is the nuclear space of all smooth, compactly supported functions on  $\mathbb{R}^d$ , and  $\mathcal{D}'$  is the dual space of  $\mathcal{D}$  with respect to the center space  $L^2(\mathbb{R}^d, dx)$ , see e.g. [2] for detail. For  $\omega \in \mathcal{D}'$  and  $\varphi \in \mathcal{D}$ , we denote by  $\langle \omega, \varphi \rangle$  the dual pairing of  $\omega$  and  $\varphi$ . Denote by  $\mathcal{C}(\mathcal{D}')$  the cylinder  $\sigma$ -algebra on  $\mathcal{D}'$ . A generalized stochastic process is a probability measure  $\mu$  on  $(\mathcal{D}', \mathcal{C}(\mathcal{D}'))$ . Thus, a generalized stochastic process is a random generalized function  $\omega \in \mathcal{D}'$ . One says that a generalized stochastic process has *independent values* if for any  $\varphi_1, \dots, \varphi_n \in \mathcal{D}$  which have mutually disjoint support, the random variables  $\langle \omega, \varphi_1 \rangle, \dots, \langle \omega, \varphi_n \rangle$  are independent. So, heuristically, we infer that, for any  $x_1, \dots, x_n \in \mathbb{R}^d$ , the random variables  $\omega(x_1), \dots, \omega(x_n)$  are independent. In the case where  $d = 1$ , one can (at least heuristically) interpret  $\omega(t)$  as the time  $t$  derivative of a classical stochastic process  $X = (X(t))_{t \in \mathbb{R}}$  with independent increments, so that, for  $t \geq 0$ ,  $X(t) = \int_0^t \omega(s) ds$ .

If a generalized stochastic process with independent values,  $\mu$ , has the property that the measure  $\mu$  remains invariant under each transformation  $x \mapsto x + a$  ( $a \in \mathbb{R}^d$ ) of the underlying space, then one calls  $\mu$  a *Lévy process* (which is, for  $d = 1$ , the time derivative of a classical Lévy process.) So, below, for a certain class of generalized stochastic processes with independent values, we will construct an orthogonal decomposition of the space  $L^2(\mathcal{D}', \mu)$ , which, in the case of a classical Lévy process, will be exactly the Nualart–Schoutens decomposition from [9]. This paper will also extend the results of [8] for generalized stochastic processes being Lévy processes.

## 2. PRELIMINARIES

We start by briefly recalling some results from [5]. Assume that, for each  $x \in \mathbb{R}^d$ ,  $\sigma(x, ds)$  is a probability measure on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ . We also assume that, for each  $\Delta \in \mathcal{B}(\mathbb{R})$ ,  $\mathbb{R}^d \ni x \mapsto \sigma(x, \Delta)$  is a measurable mapping. Hence, we can define a  $\sigma$ -finite measure  $dx \sigma(x, ds)$  on  $(\mathbb{R}^d \times \mathbb{R}, \mathcal{B}(\mathbb{R}^d \times \mathbb{R}))$ . Let  $\mathcal{B}_0(\mathbb{R}^d)$  denote the collection of all sets  $\Lambda \in \mathcal{B}(\mathbb{R}^d)$  which are bounded. We will additionally assume that, for each  $\Lambda \in \mathcal{B}_0(\mathbb{R}^d)$ , there exists  $C_\Lambda > 0$  such that

$$(2.1) \quad \int_{\mathbb{R}} |s|^n \sigma(x, ds) \leq C_\Lambda^n n!, \quad n \in \mathbb{N},$$

for all  $x \in \Lambda$ . We fix the Hilbert space  $H = L^2(\mathbb{R}^d \times \mathbb{R}, dx \sigma(x, ds))$ . We denote by  $\mathcal{F}(H) = \bigoplus_{n=0}^\infty H^{\odot n} n!$  the symmetric Fock space over  $H$ . Here  $\odot$  denotes symmetric tensor product. We denote by  $\mathfrak{D}$  the subset of  $\mathcal{F}(H)$  which consists of all finite vectors  $f = (f^{(0)}, f^{(1)}, \dots, f^{(n)}, 0, 0, \dots)$ , where each  $f^{(k)}$  is a symmet-

ric function on  $(\mathbb{R}^d \times \mathbb{R})^k$  which is obtained as the symmetrization of a finite sum of functions of the form

$$g^{(k)}(x_1, s_1, \dots, x_k, s_k) = \phi(x_1, \dots, x_k) s_1^{i_1} \dots s_k^{i_k},$$

where  $\phi \in \mathcal{D}^{\otimes k} = C_0^\infty((\mathbb{R}^d)^k)$  and  $i_1, \dots, i_k \in \mathbb{Z}_+ = \{0, 1, 2, \dots\}$ . For each  $\varphi \in \mathcal{D}$ , we define an operator  $A(\varphi)$  in  $\mathcal{F}(H)$  with domain  $\mathfrak{D}$  by

$$(2.2) \quad A(\varphi) := a^+(\varphi \otimes m_0) + a^-(\varphi \otimes m_0) + a^0(\varphi \otimes m_1).$$

Here and below, for  $i \in \mathbb{Z}_+ := \{0, 1, 2, \dots\}$ ,

$$(\varphi \otimes m_i)(x, s) := \varphi(x) s^i;$$

$a^+(\varphi \otimes m_i)$  is the creation operator corresponding to  $\varphi \otimes m_i$ ,

$$a^+(\varphi \otimes m_i) f^{(k)} = f^{(k)} \odot (\varphi \otimes m_i), \quad f^{(k)} \in H^{\odot k};$$

$a^-(\varphi \otimes m_i)$  is the corresponding annihilation operator,

$$a^-(\varphi \otimes m_i) f^{(k)} = k \int_{\mathbb{R}^d \times \mathbb{R}} dy \sigma(y, du) \varphi(y) u^i f^{(k)}(y, u, \cdot);$$

and  $a^0(\varphi \otimes m_i)$  is the neutral operator corresponding to  $\varphi \otimes m_i$ ,

$$\begin{aligned} (a^0(\varphi \otimes m_i) f^{(k)})(x_1, s_1, \dots, x_k, s_k) \\ = (\varphi(x_1) s_1^i + \dots + \varphi(x_k) s_k^i) f^{(k)}(x_1, s_1, \dots, x_k, s_k). \end{aligned}$$

Note that  $A(\varphi)$  maps  $\mathfrak{D}$  into itself, and it is a symmetric operator in  $\mathcal{F}(H)$ .

**THEOREM 2.1.** *For each  $\varphi \in \mathcal{D}$ , the operator  $A(\varphi)$  is essentially self-adjoint on  $\mathfrak{D}$ . Furthermore, there exists a unique probability measure  $\mu$  on  $\mathcal{D}'$  such that the linear operator  $I : \mathcal{F}(H) \rightarrow L^2(\mathcal{D}', \mu)$  given through  $I\Omega = 1$ ,  $\Omega$  being the vacuum vector  $(1, 0, 0, \dots)$ , and*

$$I(A(\varphi_1) \dots A(\varphi_n) \Omega) = \langle \omega, \varphi_1 \rangle \dots \langle \omega, \varphi_n \rangle,$$

*is a unitary operator. The Fourier transform of the measure  $\mu$  is given by*

$$(2.3) \quad \int_{\mathcal{D}'} e^{i\langle \varphi, \omega \rangle} \mu(d\omega) = \exp \left[ -\frac{1}{2} \int_{\mathbb{R}^d} dx \sigma(x, \{0\}) \varphi(x)^2 + \int_{\mathbb{R}^d} dx \int_{\mathbb{R}^*} \sigma(x, ds) \frac{1}{s^2} (e^{i\varphi(x)s} - i\varphi(x)s - 1) \right],$$

*where  $\mathbb{R}^* := \mathbb{R} \setminus \{0\}$ . In particular,  $\mu$  is a generalized stochastic process with independent values.*

Note that, if the measure  $\sigma(ds) = \sigma(x, ds)$  is the same for all  $x \in \mathbb{R}^d$ , then  $\mu$  is a Lévy process.

### 3. AN ORTHOGONAL DECOMPOSITION OF A FOCK SPACE

We will now discuss an orthogonal decomposition of a general symmetric Fock space. This decomposition generalizes the well-known basis of occupation numbers in the Fock space, see e.g. [2].

In this section, we will denote by  $H$  any real separable Hilbert space. Let  $(H_k)_{k=0}^\infty$  be a sequence of closed subspaces of  $H$  such that  $H = \bigoplus_{k=0}^\infty H_k$ . Let  $n \geq 2$ . Then clearly

$$(3.1) \quad \begin{aligned} H^{\otimes n} &= \left( \bigoplus_{k_1=0}^\infty H_{k_1} \right) \otimes \left( \bigoplus_{k_2=0}^\infty H_{k_2} \right) \otimes \dots \otimes \left( \bigoplus_{k_n=0}^\infty H_{k_n} \right) \\ &= \bigoplus_{(k_1, k_2, \dots, k_n) \in \mathbb{Z}_+^n} H_{k_1} \otimes H_{k_2} \otimes \dots \otimes H_{k_n}. \end{aligned}$$

Denote by  $\text{Sym}_n$  the orthogonal projection of  $H^{\otimes n}$  onto  $H^{\odot n}$ . Recall that, for any  $f_1, f_2, \dots, f_n \in H$ ,

$$(3.2) \quad f_1 \odot \dots \odot f_n = \text{Sym}_n f_1 \otimes \dots \otimes f_n = \frac{1}{n!} \sum_{\sigma \in S_n} f_{\sigma(1)} \otimes \dots \otimes f_{\sigma(n)}.$$

(Here,  $S_n$  denotes the symmetric group of order  $n$ .) For each  $(k_1, k_2, \dots, k_n) \in \mathbb{Z}_+^n$ , let us assume that  $H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n}$  denote the Hilbert space  $\text{Sym}_n(H_{k_1} \otimes H_{k_2} \otimes \dots \otimes H_{k_n})$ , i.e., the space of all  $\text{Sym}_n$ -projections of elements of  $H_{k_1} \otimes H_{k_2} \otimes \dots \otimes H_{k_n}$ .

Assume that  $(k_1, k_2, \dots, k_n) \in \mathbb{Z}_+^n$  and  $(l_1, l_2, \dots, l_n) \in \mathbb{Z}_+^n$  are such that there exists a permutation  $\sigma \in S_n$  such that

$$(3.3) \quad (k_1, k_2, \dots, k_n) = (l_{\sigma(1)}, l_{\sigma(2)}, \dots, l_{\sigma(n)}).$$

Then

$$(3.4) \quad H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n} = H_{l_1} \odot H_{l_2} \odot \dots \odot H_{l_n}.$$

Indeed, take any  $f_1 \in H_{l_1}, f_2 \in H_{l_2}, \dots, f_n \in H_{l_n}$ . Then

$$(3.5) \quad f_1 \odot f_2 \odot \dots \odot f_n = f_{\sigma(1)} \odot f_{\sigma(2)} \odot \dots \odot f_{\sigma(n)}.$$

We have  $f_{\sigma(i)} \in H_{l_{\sigma(i)}} = H_{k_i}$ . Therefore, the vector in (3.5) belongs to  $H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n}$ . Since the set of all vectors of the form  $f_1 \odot f_2 \odot \dots \odot f_n$  with  $f_i \in H_{l_i}$  is total in  $H_{l_1} \odot H_{l_2} \odot \dots \odot H_{l_n}$ , we conclude that

$$H_{l_1} \odot H_{l_2} \odot \dots \odot H_{l_n} \subset H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n}.$$

By inverting the argument, we obtain the inverse conclusion, and so formula (3.4) holds.

If no permutation  $\sigma \in S_n$  exists which satisfies (3.3), then

$$(3.6) \quad H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n} \perp H_{l_1} \odot H_{l_2} \odot \dots \odot H_{l_n}.$$

Indeed, take any  $f_i \in H_{k_i}, g_i \in H_{l_i}, i = 1, 2, \dots, n$ . Then, since  $\text{Sym}_n$  is an orthogonal projection,

$$\begin{aligned} & (f_1 \odot f_2 \odot \dots \odot f_n, g_1 \odot g_2 \odot \dots \odot g_n)_{H^{\odot n}} \\ &= (\text{Sym}_n(f_1 \otimes f_2 \otimes \dots \otimes f_n), g_1 \otimes g_2 \otimes \dots \otimes g_n)_{H^{\otimes n}} \\ &= \frac{1}{n!} \sum_{\sigma \in S_n} \prod_{i=1}^n (f_{\sigma(i)}, g_i)_H = \frac{1}{n!} \sum_{\sigma \in S_n} \prod_{i=1}^n (f_i, g_{\sigma(i)})_H = 0. \end{aligned}$$

Since the vectors of the form  $f_1 \odot f_2 \odot \dots \odot f_n$  with  $f_i \in H_{k_i}$  and  $g_1 \odot g_2 \odot \dots \odot g_n$  with  $g_i \in H_{l_i}$  form a total set in  $H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n}$  and  $H_{l_1} \odot H_{l_2} \odot \dots \odot H_{l_n}$ , respectively, we get (3.6).

By (3.1), the closed linear span of the spaces  $H_{k_1} \odot H_{k_2} \odot \dots \odot H_{k_n}$  with  $(k_1, k_2, \dots, k_n) \in \mathbb{Z}_+^n$  coincides with  $H^{\odot n}$ . Hence, by (3.4) and (3.6), we get the orthogonal decomposition

$$(3.7) \quad H^{\odot n} = \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty, |\alpha|=n} H_0^{\odot \alpha_0} \odot H_1^{\odot \alpha_1} \odot H_2^{\odot \alpha_2} \odot \dots$$

Here  $\mathbb{Z}_{+,0}^\infty$  denotes the set of indices  $\alpha = (\alpha_0, \alpha_1, \alpha_2, \dots)$  such that all  $\alpha_i \in \mathbb{Z}_+$  and  $|\alpha| := \alpha_0 + \alpha_1 + \alpha_2 + \dots < \infty$ . Hence, by (3.7), we get the following

LEMMA 3.1. *We have the orthogonal decomposition of the symmetric Fock space  $\mathcal{F}(H) = \bigoplus_{n=0}^\infty H^{\odot n} n!$ , i.e.,*

$$(3.8) \quad \mathcal{F}(H) = \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} (H_0^{\odot \alpha_0} \odot H_1^{\odot \alpha_1} \odot H_2^{\odot \alpha_2} \dots) |\alpha|!$$

Next, we have

LEMMA 3.2. *Let  $\alpha \in \mathbb{Z}_{+,0}^\infty$ . Then*

$$(3.9) \quad \begin{aligned} \text{Sym}_{|\alpha|} : (H_0^{\odot \alpha_0} \otimes H_1^{\odot \alpha_1} \otimes H_2^{\odot \alpha_2} \otimes \dots) \alpha_0! \alpha_1! \alpha_2! \dots \\ \rightarrow (H_0^{\odot \alpha_0} \odot H_1^{\odot \alpha_1} \odot H_2^{\odot \alpha_2} \odot \dots) |\alpha|! \end{aligned}$$

is a unitary operator.

Proof. We start the proof with the following well-known observation. Let  $k, l \geq 1, n := k + l$ . Then  $\text{Sym}_n = \text{Sym}_n(\text{Sym}_k \otimes \text{Sym}_l)$ . Hence, for any  $\alpha \in$

$\mathbb{Z}_{+,0}^\infty$ ,  $|\alpha| = n$ , we get  $\text{Sym}_n = \text{Sym}_n(\text{Sym}_{\alpha_0} \otimes \text{Sym}_{\alpha_1} \otimes \text{Sym}_{\alpha_2} \otimes \dots)$ . Therefore, we have the following equality of subspaces of  $H^{\otimes n}$ :

$$\begin{aligned} & H_0^{\odot\alpha_0} \odot H_1^{\odot\alpha_1} \odot H_2^{\odot\alpha_2} \odot \dots \\ &= \text{Sym}_n(H_0^{\otimes\alpha_0} \otimes H_1^{\otimes\alpha_1} \otimes H_2^{\otimes\alpha_2} \otimes \dots) \\ &= \text{Sym}_n(\text{Sym}_{\alpha_0} \otimes \text{Sym}_{\alpha_1} \otimes \text{Sym}_{\alpha_2} \otimes \dots)(H_0^{\otimes\alpha_0} \otimes H_1^{\otimes\alpha_1} \otimes H_2^{\otimes\alpha_2} \otimes \dots) \\ &= \text{Sym}_n(H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes H_2^{\odot\alpha_2} \otimes \dots). \end{aligned}$$

This shows that the image of the operator  $\text{Sym}_n$  in (3.9) is the whole space  $H_0^{\odot\alpha_0} \odot H_1^{\odot\alpha_1} \odot H_2^{\odot\alpha_2} \odot \dots n!$ . Hence, we only need to prove that this operator is an isometry.

Fix any  $f_i, g_i \in H_i$  with  $i \in \mathbb{Z}_+$ , and any  $\alpha \in \mathbb{Z}_{+,0}^\infty$ . Then, by (3.2),

$$\begin{aligned} & (\text{Sym}_n(f_0^{\otimes\alpha_0} \otimes f_1^{\otimes\alpha_1} \otimes f_2^{\otimes\alpha_2} \otimes \dots), \text{Sym}_n(g_0^{\otimes\alpha_0} \otimes g_1^{\otimes\alpha_1} \otimes g_2^{\otimes\alpha_2} \otimes \dots))_{H^{\odot n}} \\ &= (\text{Sym}_n(f_0^{\otimes\alpha_0} \otimes f_1^{\otimes\alpha_1} \otimes f_2^{\otimes\alpha_2} \otimes \dots), g_0^{\otimes\alpha_0} \otimes g_1^{\otimes\alpha_1} \otimes g_2^{\otimes\alpha_2} \otimes \dots)_{H^{\otimes n}} \\ &= \frac{1}{n!} \sum_{\sigma_0 \in S_{\alpha_0}} (f_0, g_0)_{H_0}^{\alpha_0} \cdot \sum_{\sigma_1 \in S_{\alpha_1}} (f_1, g_1)_{H_1}^{\alpha_1} \dots \\ &= \frac{1}{n!} (f_0^{\otimes\alpha_0}, g_0^{\otimes\alpha_0})_{H_0^{\odot\alpha_0}} \alpha_0! (f_1^{\otimes\alpha_1}, g_1^{\otimes\alpha_1})_{H_1^{\odot\alpha_1}} \alpha_1! \dots \\ &= \frac{1}{n!} (f_0^{\otimes\alpha_0} \otimes f_1^{\otimes\alpha_1} \otimes \dots, g_0^{\otimes\alpha_0} \otimes g_1^{\otimes\alpha_1} \otimes \dots)_{H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes \dots} \alpha_0! \alpha_1! \dots \end{aligned}$$

Since the set of all vectors of the form  $f_i^{\otimes\alpha_i}$  with  $f_i \in H_i$  is a total subset of  $H_i^{\odot\alpha_i}$ , we conclude that the operator in (3.9) is indeed an isometry. ■

We define the symmetrization operator

$$(3.10) \quad \text{Sym} : \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} (H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes H_2^{\odot\alpha_2} \otimes \dots) \alpha_0! \alpha_1! \alpha_2! \dots \rightarrow \mathcal{F}(H)$$

so that the restriction of  $\text{Sym}$  to each space

$$(H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes H_2^{\odot\alpha_2} \otimes \dots) \alpha_0! \alpha_1! \alpha_2! \dots$$

is equal to  $\text{Sym}_{|\alpha|}$ . By Lemmas 3.1 and 3.2, we get

LEMMA 3.3. *The symmetrization operator  $\text{Sym}$  is a unitary operator.*

REMARK 3.1. *Let us assume that each Hilbert space  $H_k$  is one-dimensional and in each  $H_k$  we fix a vector  $e_k \in H_k$  such that  $\|e_k\| = 1$ . Thus,  $(e_k)_{k=0}^\infty$  is an orthonormal basis of  $H$ . By Lemma 3.3, the set of the vectors*

$$((\alpha_0! \alpha_1! \alpha_2! \dots)^{-1/2} e_0^{\otimes\alpha_0} \odot e_1^{\otimes\alpha_1} \odot e_2^{\otimes\alpha_2} \odot \dots)_{\alpha \in \mathbb{Z}_{+,0}^\infty}$$

*is an orthonormal basis of  $\mathcal{F}(H)$ . This basis is called a basis of occupation numbers.*

4. AN ORTHOGONAL DECOMPOSITION OF  $L^2(\mathcal{D}', \mu)$

We want to apply the general result about the orthogonal decomposition of the Fock space to the case of  $\mathcal{F}(H)$ , where  $H = L^2(\mathbb{R}^d \times \mathbb{R}, dx \sigma(x, ds))$ . We note that, by (2.1), for each  $x \in \mathbb{R}^d$ , the set of polynomials is dense in  $L^2(\mathbb{R}, \sigma(x, ds))$ . We denote by  $(q^{(n)}(x, s))_{n \geq 0}$  the sequence of monic polynomials which are orthogonal with respect to the measure  $\sigma(x, ds)$ . These polynomials satisfy the following recursive formula:

$$(4.1) \quad \begin{aligned} sq^{(n)}(x, s) &= q^{(n+1)}(x, s) + b_n(x)q^{(n)}(x, s) + a_n(x)q^{(n-1)}(x, s), \quad n \geq 1, \\ sq^{(0)}(x, s) &= q^{(1)}(x, s) + b_0(x) \end{aligned}$$

with some  $b_n(x) \in \mathbb{R}$  and  $a_n(x) > 0$ . (Note that if the support of  $\sigma(x, ds)$  consists of  $k < \infty$  points, then, for  $n \geq k$ , we set  $q^{(n)}(x, s) = 0$ ,  $a_n(x) = 0$  with  $b_n(x) \in \mathbb{R}$  being arbitrary.)

From now on, we will assume that the following condition is satisfied:

(A) For each  $n \in \mathbb{N}$ , the function  $a_n(x)$  from (4.1) is locally bounded on  $\mathbb{R}^d$ , i.e., for each  $\Lambda \in \mathcal{B}_0(\mathbb{R}^d)$ ,  $\sup_{x \in \Lambda} a_n(x) < \infty$ .

Denote by  $\mathcal{L}$  the linear space of all functions on  $\mathbb{R}^d \times \mathbb{R}$  which have the form

$$(4.2) \quad f(x, s) = \sum_{k=0}^n a_k(x)q^{(k)}(x, s),$$

where  $n \in \mathbb{N}$ ,  $a_k \in \mathcal{D}$ ,  $k = 0, 1, \dots, n$ .

LEMMA 4.1. *The space  $\mathcal{L}$  is densely embedded into  $H$ .*

Proof. Let  $f(x, s) = a(x)q^{(k)}(x, s)$ , where  $a \in \mathcal{D}$ . Let us show that  $f \in H$ . Put  $\Lambda := \text{supp}(a)$ . We have, for some  $C > 0$ ,

$$(4.3) \quad \int_{\mathbb{R}^d} \int_{\mathbb{R}} dx \sigma(x, ds) f(x, s)^2 \leq C \int_{\Lambda} dx \int_{\mathbb{R}} \sigma(x, ds) q^{(k)}(x, s)^2.$$

If  $k = 0$ , then  $q^{(0)}(x, s) = 1$ , and the right-hand side of (4.3) is evidently finite. By the theory of orthogonal polynomials (see e.g. [4])

$$(4.4) \quad \int_{\mathbb{R}} \sigma(x, ds) q^{(k)}(x, s)^2 = a_1(x)a_2(x) \dots a_k(x), \quad k \geq 1.$$

Hence we continue (4.3) and obtain

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}} dx \sigma(x, ds) f(x, s)^2 \leq C \int_{\Lambda} dx a_1(x)a_2(x) \dots a_k(x) < \infty$$

by (A). Thus,  $\mathcal{L} \subset H$ .

We now have to show that  $\mathfrak{L}$  is a dense subset of  $H$ . Let  $g \in H$  be such that  $(g, f)_H = 0$  for all  $f \in \mathfrak{L}$ . Hence for any  $a \in \mathcal{D}$  and  $k \geq 0$

$$\int_{\mathbb{R}^d} dx \int_{\mathbb{R}} \sigma(x, ds) g(x, s) a(x) q^{(k)}(x, s) = 0.$$

Fix any compact set  $\Lambda$  in  $\mathbb{R}^d$  and let  $a \in \mathcal{D}$  be such that the support of  $a$  is a subset of  $\Lambda$ . Then,

$$\int_{\mathbb{R}^d} dx a(x) \left( \int_{\mathbb{R}} \sigma(x, ds) g(x, s) q^{(k)}(x, s) \right) = 0.$$

Hence

$$(4.5) \quad \int_{\Lambda} dx a(x) \left( \int_{\mathbb{R}} \sigma(x, ds) g(x, s) q^{(k)}(x, s) \right) = 0.$$

We state that the function

$$\Lambda \ni x \mapsto \int_{\mathbb{R}} \sigma(x, ds) g(x, s) q^{(k)}(x, s)$$

belongs to  $L^2(\Lambda, dx)$ . Indeed, if  $k = 0$ , then  $q^{(0)}(x, s) = 1$ , and this statement evidently follows from Cauchy's inequality. Assume that  $k \geq 1$ . Then, by Cauchy's inequality, (4.3), and condition (A),

$$\begin{aligned} & \int_{\Lambda} dx \left( \int_{\mathbb{R}} \sigma(x, ds) g(x, s) q^{(k)}(x, s) \right)^2 \\ & \leq \int_{\Lambda} dx \int_{\mathbb{R}} \sigma(x, ds_1) g(x, s_1)^2 \int_{\mathbb{R}} \sigma(x, ds_2) q^{(k)}(x, s_2)^2 \\ & = \int_{\Lambda} dx \int_{\mathbb{R}} \sigma(x, ds) g(x, s)^2 a_1(x) a_2(x) \dots a_k(x) \\ & \leq \left( \prod_{i=1}^k \sup_{x \in \Lambda} a_i(x) \right) \int_{\Lambda} dx \int_{\mathbb{R}} \sigma(x, ds) g(x, s)^2 < \infty. \end{aligned}$$

Since the set of all functions  $a \in \mathcal{D}$  with support in  $\Lambda$  is dense in  $L^2(\Lambda, dx)$ , we therefore conclude from (4.5) that, for  $dx$ -a.a.  $x \in \Lambda$ ,

$$(4.6) \quad \int_{\mathbb{R}} \sigma(x, ds) g(x, s) q^{(k)}(x, s) = 0 \quad \text{for all } k \geq 0.$$

Since  $g \in H$ , we infer that, for  $dx$ -a.a.  $x \in \mathbb{R}^d$ ,  $g(x, \cdot) \in L^2(\mathbb{R}, \sigma(x, ds))$ . Since  $\{q^{(k)}(x, \cdot)\}_{k=0}^{\infty}$  form an orthogonal basis in  $L^2(\mathbb{R}, \sigma(x, ds))$ , we conclude from (4.6) that, for  $dx$ -a.a.  $x \in \mathbb{R}^d$ ,  $g(x, s) = 0$  for  $\sigma(x, ds)$ -a.a.  $s \in \mathbb{R}$ . From here, we easily conclude that  $g = 0$  as an element of  $H$ . Hence  $\mathfrak{L}$  is indeed dense in  $H$ . ■



For each  $n \in \mathbb{Z}_+$ , we define

$$\mathfrak{L}_n := \{g_n(x, s) = f(x) q^{(n)}(x, s) \mid f \in \mathcal{D}\}.$$

We have  $\mathfrak{L}_n \subset \mathfrak{L}$ , and so the linear span of the  $\mathfrak{L}_n$  spaces coincides with  $\mathfrak{L}$ . For any  $g_n(x, s) = f_n(x) q^{(n)}(x, s) \in \mathfrak{L}_n$  and  $g_m(x, s) = f_m(x) q^{(m)}(x, s) \in \mathfrak{L}_m$ ,  $n, m \in \mathbb{Z}_+$ , we have

$$\begin{aligned} (4.7) \quad (g_n, g_m)_H &= \int_{\mathbb{R}^d \times \mathbb{R}} g_n(x, s) g_m(x, s) dx \sigma(x, ds) \\ &= \int_{\mathbb{R}^d} f_n(x) f_m(x) \left( \int_{\mathbb{R}} q^{(n)}(x, s) q^{(m)}(x, s) \sigma(x, ds) \right) dx. \end{aligned}$$

Hence, if  $n \neq m$ , then

$$(g_n, g_m)_H = 0,$$

which implies that the linear spaces  $\{\mathfrak{L}_n\}_{n=0}^\infty$  are mutually orthogonal in  $H$ . Denote by  $H_n$  the closure of  $\mathfrak{L}_n$  in  $H$ . Then, by Lemma 4.1,  $H = \bigoplus_{n=0}^\infty H_n$ .

By (4.7), setting  $n = m$ , we get

$$(4.8) \quad \|g_n\|_{H_n}^2 = \int_{\mathbb{R}^d} f_n^2(x) \left( \int_{\mathbb{R}} q^{(n)}(x, s)^2 \sigma(x, ds) \right) dx = \int_{\mathbb{R}^d} f_n^2(x) \rho_n(dx),$$

where

$$\rho_n(dx) = \left( \int_{\mathbb{R}} q^{(n)}(x, s)^2 \sigma(x, ds) \right) dx$$

is a measure on  $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ . Consider a linear operator

$$\mathcal{D} \ni f_n \mapsto (J_n f_n)(x, s) := f_n(x) q^{(n)}(x, s) \in \mathfrak{L}_n.$$

The image of  $J_n$  is clearly the whole  $\mathfrak{L}_n$ . Now,  $\mathfrak{L}_n$  is dense in  $H_n$ , while  $\mathcal{D}$  is evidently dense in  $L^2(\mathbb{R}^d, \rho_n(dx))$ . By (4.8), for each  $f_n \in \mathcal{D}$ ,

$$\|J_n f_n\|_{H_n} = \|f_n\|_{L^2(\mathbb{R}^d, \rho_n(dx))}.$$

Therefore, we can extend the operator  $J_n$  by continuity to a unitary operator

$$(4.9) \quad J_n : L^2(\mathbb{R}^d, \rho_n(dx)) \rightarrow H_n.$$

In particular,

$$H_n = \{f_n(x) q^{(n)}(x, s) \mid f_n \in L^2(\mathbb{R}^d, \rho_n(dx))\}.$$

Therefore, for each  $k \geq 2$ ,

$$\begin{aligned} H_n^{\otimes k} &= \{f_n^{(k)}(x_1, \dots, x_k) q^{(n)}(x_1, s_1) \dots q^{(n)}(x_k, s_k) \mid \\ &\quad f_n^{(k)} \in L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k} = L^2((\mathbb{R}^d)^k, \rho_n(dx_1) \dots \rho_n(dx_k))\}. \end{aligned}$$

Since the operator  $J_n$  in (4.9) is unitary, we infer that the operator

$$J_n^{\otimes k} : L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k} \rightarrow H_n^{\otimes k}$$

is also unitary. The restriction of  $J_n^{\otimes k}$  to  $L^2(\mathbb{R}^d, \rho_n(dx))^{\odot k}$  is a unitary operator

$$(4.10) \quad J_n^{\otimes k} : L^2(\mathbb{R}^d, \rho_n(dx))^{\odot k} \rightarrow H_n^{\odot k}.$$

Indeed, take any  $f_n \in L^2(\mathbb{R}^d, \rho_n(dx))$ . Then  $f_n^{\otimes k} \in L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k}$  and the set of all such vectors is total in  $L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k}$ . Now, by the definition of  $J_n^{\otimes k}$ , we get

$$J_n^{\otimes k} f_n^{\otimes k} = (J_n f_n)^{\otimes k} \in H_n^{\otimes k},$$

and furthermore the set of all vectors of the form  $(J_n f_n)^{\otimes k}$  is total in  $H_n^{\otimes k}$ . Hence, the statement follows.

For any  $f_n^{(k)} \in L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k}$ ,

$$(J_n^{\otimes k} f_n^{(k)})(x_1, s_1, \dots, x_k, s_k) = f_n^{(k)}(x_1, \dots, x_k) q^{(n)}(x_1, s_1) \dots q^{(n)}(x_k, s_k).$$

Hence, the unitary operator (4.10) acts as follows:

$$\begin{aligned} L^2(\mathbb{R}^d, \rho_n(dx))^{\odot k} &\ni f_n^{(k)}(x_1, \dots, x_k) \\ \mapsto (J_n^{\otimes k} f_n^{(k)})(x_1, s_1, \dots, x_k, s_k) &= f_n^{(k)}(x_1, \dots, x_k) q^{(n)}(x_1, s_1) \dots q^{(n)}(x_k, s_k). \end{aligned}$$

Thus, each function  $g_n^{(k)} \in H_n^{\odot k}$  has a representation

$$g_n^{(k)}(x_1, s_1, \dots, x_k, s_k) = f_n^{(k)}(x_1, \dots, x_k) q^{(n)}(x_1, s_1) \dots q^{(n)}(x_k, s_k),$$

where  $f_n^{(k)} \in L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k}$  and  $\|g_n^{(k)}\|_{H_n^{\odot k}} = \|f_n^{(k)}\|_{L^2(\mathbb{R}^d, \rho_n(dx))^{\otimes k}}$ .

For each  $\alpha \in \mathbb{Z}_{+,0}^\infty$ , we consider the Hilbert space

$$(4.11) \quad L_\alpha^2((\mathbb{R}^d)^{|\alpha|}) := L^2(\mathbb{R}^d, \rho_0(dx))^{\odot \alpha_0} \otimes L^2(\mathbb{R}^d, \rho_1(dx))^{\odot \alpha_1} \otimes \dots$$

We now define a unitary operator

$$J_\alpha : L_\alpha^2((\mathbb{R}^d)^{|\alpha|}) \rightarrow H_0^{\odot \alpha_0} \otimes H_1^{\odot \alpha_1} \otimes \dots,$$

where

$$J_\alpha = J_0^{\otimes \alpha_0} \otimes J_1^{\otimes \alpha_1} \otimes \dots$$

We evidently have, for each  $f_\alpha \in L_\alpha^2((\mathbb{R}^d)^{|\alpha|})$ ,

$$\begin{aligned} (J_\alpha f_\alpha)(x_1, s_1, x_2, s_2, \dots, x_{|\alpha|}, s_{|\alpha|}) \\ = f_\alpha(x_1, x_2, \dots, x_{|\alpha|}) q^{(0)}(x_1, s_1) \dots q^{(0)}(x_{\alpha_0}, s_{\alpha_0}) \\ \times q^{(1)}(x_{\alpha_0+1}, s_{\alpha_0+1}) \dots q^{(1)}(x_{\alpha_0+\alpha_1}, s_{\alpha_0+\alpha_1}) \dots \end{aligned}$$

For each  $\alpha \in \mathbb{Z}_{+,0}^\infty$ , we define a Hilbert space

$$\mathcal{G}_\alpha := L_\alpha^2((\mathbb{R}^d)^{|\alpha|})\alpha_0!\alpha_1!\dots$$

The  $J_\alpha$  is evidently a unitary operator

$$J_\alpha : \mathcal{G}_\alpha \rightarrow (H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes \dots)\alpha_0!\alpha_1!\dots$$

Put  $\mathcal{G} := \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} \mathcal{G}_\alpha$ . Hence, we can construct a unitary operator

$$J : \mathcal{G} \rightarrow \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} (H_0^{\odot\alpha_0} \otimes H_1^{\odot\alpha_1} \otimes \dots)\alpha_0!\alpha_1!\dots$$

by setting  $J := \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} J_\alpha$ . By Lemma 3.3, we get a unitary operator  $\mathcal{R} : \mathcal{G} \rightarrow \mathcal{F}(H)$  by setting  $\mathcal{R} := \text{Sym } J$ . Thus, by Theorem 2.1, we get

**THEOREM 4.1.** *Let condition (A) be satisfied. We have a unitary isomorphism  $\mathcal{K} : \mathcal{G} \rightarrow L^2(\mathcal{D}', \mu)$  given by  $\mathcal{K} := I\mathcal{R}$ , where the unitary operator  $I : \mathcal{F}(H) \rightarrow L^2(\mathcal{D}', \mu)$  is from Theorem 2.1.*

**5. THE UNITARY ISOMORPHISM  $\mathcal{K}$  THROUGH MULTIPLE STOCHASTIC INTEGRALS**

We will now give an interpretation of the unitary isomorphism  $\mathcal{K}$  in terms of multiple stochastic integrals. We will only present a sketch of the proof, omitting some technical details.

Let us recall the operators  $A(\varphi)$  in  $\mathcal{F}(H)$  defined by (2.2). Now, for each  $k \in \mathbb{N}$ , we define operators

$$(5.1) \quad A^{(k)}(\varphi) := a^+(\varphi \otimes m_{k-1}) + a^0(\varphi \otimes m_k) + a^-(\varphi \otimes m_{k-1}).$$

In particular,  $A^{(1)}(\varphi) = A(\varphi)$ . The operator  $A^{(k)}(\varphi)$  being symmetric, we denote by  $A^{(k)}(\varphi)^\sim$  the closure of  $A^{(k)}(\varphi)$ . For each  $k \in \mathbb{N}$  and  $\varphi \in \mathcal{D}$ , we define  $Y^{(k-1)}(\varphi) := I(\varphi \otimes m_{k-1})$ . It can be shown that, for each  $k \in \mathbb{N}$ ,  $IA^{(k)}(\varphi)^\sim I^{-1}$  is the operator of multiplication by the function  $Y^{(k-1)}$ .

Suppose, for a moment, that the measures  $\sigma(x, ds)$  do not depend on  $x \in \mathbb{R}^d$ . For a fixed  $\varphi \in \mathcal{D}$ , let us orthogonalize in  $L^2(\mathcal{D}', \mu)$  the functions  $(Y^{(k)}(\varphi))_{k=0}^\infty$ . This is of course equivalent to the orthogonalization of the monomials  $(s^k)_{k=0}^\infty$  in  $L^2(\mathbb{R}, \sigma)$ . Denote by  $(q^{(k)})_{k=0}^\infty$  the system of monic orthogonal polynomials with respect to the measure  $\sigma$ . Let us put  $(\varphi \otimes q^{(k)})(x, s) := \varphi(x)q^{(k)}(s)$ . Thus, the random variables

$$Z^{(k)}(\varphi) := I(\varphi \otimes q^{(k)}), \quad k \in \mathbb{Z}_+,$$

appear as a result of the orthogonalization of  $(Y^{(k)}(\varphi))_{k=0}^\infty$ . Since  $q^{(0)}(s) = 1$ , we have

$$Z^{(0)}(\varphi) = Y^{(0)}(\varphi) = \langle \cdot, \varphi \rangle.$$

For each  $k \geq 1$ , we have a representation of  $q^{(k)}(s)$  as follows:

$$q^{(k)}(s) = \sum_{i=0}^k b_i^{(k)} s^i.$$

Thus,

$$Z^{(k)}(\varphi) = I(\varphi \otimes q^{(k)}) = \sum_{i=0}^k b_i^{(k)} I(\varphi \otimes m_i) = \sum_{i=0}^k b_i^{(k)} Y^{(i)}(\varphi).$$

Hence, under  $I^{-1}$ , the image of the operator of multiplication by  $Z^{(k)}(\varphi)$  is the operator

$$\begin{aligned} R^{(k)}(\varphi) &:= \sum_{i=0}^k b_i^{(k)} (a^+(\varphi \otimes m_i) + a^-(\varphi \otimes m_i) + a^0(\varphi \otimes m_{i+1})) \\ &= a^+(\varphi \otimes q^{(k)}) + a^-(\varphi \otimes q^{(k)}) + a^0(\varphi \otimes \rho^{(k)}), \end{aligned}$$

where  $\rho^{(k)}(s) := sq^{(k)}(s)$ .

Let us now consider the general case, i.e., the case where the measure  $\sigma(x, ds)$  does depend on  $x \in \mathbb{R}^d$ . We are using the monic polynomials  $(q^{(k)}(x, \cdot))_{k=0}^{\infty}$  which are orthogonal with respect to the measure  $\sigma(x, ds)$ . We have

$$q^{(k)}(x, s) = \sum_{i=0}^k b_i^{(k)}(x) s^i.$$

We define

$$Z^{(k)}(\varphi) := I(\varphi q^{(k)}) = \sum_{i=0}^k Y^{(i)}(\varphi b_i^{(k)}),$$

where  $(\varphi q^{(k)})(x, s) := \varphi(x) q^{(k)}(x, s)$ . Hence, under  $I^{-1}$ , the image of the operator of multiplication by  $Z^{(k)}(\varphi)$  is the operator

$$\begin{aligned} R^{(k)}(\varphi) &:= \sum_{i=0}^k \left( a^+((\varphi b_i^{(k)}) \otimes m_i) + a^-((\varphi b_i^{(k)}) \otimes m_i) + a^0((\varphi b_i^{(k)}) \otimes m_{i+1}) \right) \\ &= a^+ \left( \left( \varphi \sum_{i=0}^k b_i^{(k)} \right) \otimes m_i \right) + a^- \left( \left( \varphi \sum_{i=0}^k b_i^{(k)} \right) \otimes m_i \right) \\ &\quad + a^0 \left( \left( \varphi \sum_{i=0}^k b_i^{(k)} \right) \otimes m_{i+1} \right) \\ &= a^+(\varphi q^{(k)}) + a^-(\varphi q^{(k)}) + a^0(\varphi \rho^{(k)}), \end{aligned}$$

where  $\rho^{(k)}(x, s) := sq^{(k)}(x, s)$ .

It is not hard to see that the above definitions and formulas can be easily extended to the case where the function  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  is just measurable, bounded, and has compact support. In particular, for each  $\Delta \in \mathcal{B}_0(\mathbb{R}^d)$ , we will use the operators  $Z^{(k)}(\Delta) := Z^{(k)}(\chi_\Delta)$ .

We will now introduce a multiple Wiener–Itô integral with respect to  $Z^{(k)}$ 's. So, we fix any  $\alpha \in \mathbb{Z}_{+,0}^\infty$ ,  $|\alpha| = n$ ,  $n \in \mathbb{N}$ . Take any  $\Delta_1, \dots, \Delta_n \in \mathcal{B}_0(\mathbb{R}^d)$ , mutually disjoint. Then we define

$$\begin{aligned} & \int_{\Delta_1 \times \Delta_2 \times \dots \times \Delta_n} dZ^{(0)}(x_1) \dots dZ^{(0)}(x_{\alpha_0}) dZ^{(1)}(x_{\alpha_0+1}) \dots dZ^{(1)}(x_{\alpha_0+\alpha_1}) \\ & \quad \times dZ^{(2)}(x_{\alpha_0+\alpha_1+1}) \dots \\ &= \int_{(\mathbb{R}^d)^n} \chi_{\Delta_1}(x_1) \chi_{\Delta_2}(x_2) \dots \chi_{\Delta_n}(x_n) dZ^{(0)}(x_1) \dots dZ^{(0)}(x_{\alpha_0}) \\ & \quad \times dZ^{(1)}(x_{\alpha_0+1}) \dots dZ^{(1)}(x_{\alpha_0+\alpha_1}) dZ^{(2)}(x_{\alpha_0+\alpha_1+1}) \dots \\ &:= Z^{(0)}(\Delta_1) \dots Z^{(0)}(\Delta_{\alpha_0}) Z^{(1)}(\Delta_{\alpha_0+1}) \dots Z^{(1)}(\Delta_{\alpha_0+\alpha_1}) Z^{(2)}(\Delta_{\alpha_0+\alpha_1+1}) \dots \end{aligned}$$

Using the fact that the sets  $\Delta_1, \dots, \Delta_n$  are mutually disjoint, we get

$$\begin{aligned} & I^{-1}(Z^{(0)}(\Delta_1) \dots Z^{(0)}(\Delta_{\alpha_0}) Z^{(1)}(\Delta_{\alpha_0+1}) \dots Z^{(1)}(\Delta_{\alpha_0+\alpha_1}) Z^{(2)}(\Delta_{\alpha_0+\alpha_1+1}) \dots) \\ &= R^{(0)}(\chi_{\Delta_1}) \dots R^{(0)}(\chi_{\Delta_{\alpha_0}}) R^{(1)}(\chi_{\Delta_{\alpha_0+1}}) \dots R^{(1)}(\chi_{\Delta_{\alpha_0+\alpha_1}}) R^{(2)}(\chi_{\Delta_{\alpha_0+\alpha_1+1}}) \dots \\ &= a^+(\chi_{\Delta_1} q^{(0)}) \dots a^+(\chi_{\Delta_{\alpha_0}} q^{(0)}) a^+(\chi_{\Delta_{\alpha_0+1}} q^{(1)}) \dots a^+(\chi_{\Delta_{\alpha_0+\alpha_1}} q^{(1)}) \\ & \quad \times a^+(\chi_{\Delta_{\alpha_0+\alpha_1+1}} q^{(2)}) \dots \Omega \\ &= (\chi_{\Delta_1} q^{(0)}) \odot \dots \odot (\chi_{\Delta_{\alpha_0}} q^{(0)}) \odot (\chi_{\Delta_{\alpha_0+1}} q^{(1)}) \odot \dots \odot (\chi_{\Delta_{\alpha_0+\alpha_1}} q^{(1)}) \\ & \quad \odot (\chi_{\Delta_{\alpha_0+\alpha_1+1}} q^{(2)}) \odot \dots \\ &= \text{Sym}_n \left( [(\chi_{\Delta_1} q^{(0)}) \odot \dots \odot (\chi_{\Delta_{\alpha_0}} q^{(0)})] \right. \\ & \quad \left. \otimes [(\chi_{\Delta_{\alpha_0+1}} q^{(1)}) \odot \dots \odot (\chi_{\Delta_{\alpha_0+\alpha_1}} q^{(1)})] \otimes \dots \right) \\ &= \text{Sym}_n \left( [(\chi_{\Delta_1} \odot \dots \odot \chi_{\Delta_{\alpha_0}})(x_1, \dots, x_{\alpha_0}) q^{(0)}(x_1, s_1) \dots q^{(0)}(x_{\alpha_0}, s_{\alpha_0})] \right. \\ & \quad \left. \otimes [(\chi_{\Delta_{\alpha_0+1}} \odot \dots \odot \chi_{\Delta_{\alpha_0+\alpha_1}})(x_{\alpha_0+1}, \dots, x_{\alpha_0+\alpha_1}) q^{(1)}(x_{\alpha_0+1}, s_{\alpha_0+1}) \right. \\ & \quad \left. \dots q^{(1)}(x_{\alpha_0+\alpha_1}, s_{\alpha_0+\alpha_1})] \otimes \dots \right) \\ &= \mathcal{R} \left( (\chi_{\Delta_1} \odot \dots \odot \chi_{\Delta_{\alpha_0}}) \otimes (\chi_{\Delta_{\alpha_0+1}} \odot \dots \odot \chi_{\Delta_{\alpha_0+\alpha_1}}) \otimes \dots \right). \end{aligned}$$

Hence

$$\begin{aligned} & Z^{(0)}(\Delta_1) \dots Z^{(0)}(\Delta_{\alpha_0}) Z^{(1)}(\Delta_{\alpha_0+1}) \dots Z^{(1)}(\Delta_{\alpha_0+\alpha_1}) Z^{(2)}(\Delta_{\alpha_0+\alpha_1+1}) \dots \\ & \quad = \mathcal{K} \left( (\chi_{\Delta_1} \odot \dots \odot \chi_{\Delta_{\alpha_0}}) \otimes (\chi_{\Delta_{\alpha_0+1}} \odot \dots \odot \chi_{\Delta_{\alpha_0+\alpha_1}}) \otimes \dots \right). \end{aligned}$$

The set of all vectors of the form

$$((\chi_{\Delta_1} \odot \dots \odot \chi_{\Delta_{\alpha_0}}) \otimes (\chi_{\Delta_{\alpha_0+1}} \odot \dots \odot \chi_{\Delta_{\alpha_0+\alpha_1}}) \otimes \dots)$$

is total in  $\mathcal{G}_\alpha$ . Therefore, by linearity and continuity, we can extend the definition of the multiple Wiener–Itô integral to the whole space  $\mathcal{G}_\alpha$ . Thus, we get, for each  $f_\alpha \in \mathcal{G}_\alpha$ ,

$$\int_{(\mathbb{R}^d)^{|\alpha|}} f_\alpha(x_1, \dots, x_{|\alpha|}) dZ^{(0)}(x_1) \dots dZ^{(0)}(x_{\alpha_0}) dZ^{(1)}(x_{\alpha_0+1}) \dots dZ^{(1)}(x_{\alpha_0+\alpha_1}) \\ \times dZ^{(2)}(x_{\alpha_0+\alpha_1+1}) \dots = \mathcal{K}f_\alpha.$$

Thus, we have the following theorem.

**THEOREM 5.1.** *The unitary isomorphism  $\mathcal{K}: \mathcal{G} \rightarrow L^2(\mathcal{D}', \mu)$  from Theorem 4.1 is given by*

$$\mathcal{G} = \bigoplus_{\alpha \in \mathbb{Z}_{+,0}^\infty} \mathcal{G}_\alpha \ni (f_\alpha)_{\alpha \in \mathbb{Z}_{+,0}^\infty} = f \mapsto \mathcal{K}f \\ = \sum_{\alpha \in \mathbb{Z}_{+,0}^\infty} \int_{(\mathbb{R}^d)^{|\alpha|}} f_\alpha(x_1, \dots, x_{|\alpha|}) dZ^{(0)}(x_1) \dots dZ^{(0)}(x_{\alpha_0}) \\ \times dZ^{(1)}(x_{\alpha_0+1}) \dots dZ^{(1)}(x_{\alpha_0+\alpha_1}) dZ^{(2)}(x_{\alpha_0+\alpha_1+1}) \dots$$

**Acknowledgments.** E.L. acknowledges the financial support of the Polish National Science Centre, grant no. Dec-2012/05/B/ST1/00626. The authors are grateful to the anonymous referee for many useful suggestions.

#### REFERENCES

- [1] M. Anshelevich, *q-Lévy processes*, J. Reine Angew. Math. 576 (2004), pp. 81–207.
- [2] Y. M. Berezansky and Y. G. Kondratiev, *Spectral Methods in Infinite-Dimensional Analysis. Vol. 1*, Kluwer Academic Publishers, Dordrecht 1995.
- [3] M. Bożejko and E. Lytvynov, *Meixner class of non-commutative generalized stochastic processes with freely independent values. I. A characterization*, Comm. Math. Phys. 292 (2009), pp. 99–129.
- [4] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Math. Appl., Vol. 13, Gordon and Breach Science Publishers, New York–London–Paris 1978.
- [5] S. Das, *Orthogonal Decompositions for Generalized Stochastic Processes with Independent Values*, PhD thesis, Swansea 2013.
- [6] G. Di Nunno, B. Øksendal, and F. Proske, *Malliavin Calculus for Lévy Processes with Applications to Finance*, Universitext, Springer, Berlin 2009.
- [7] J. Lin, *Chaotic and predictable representations for multidimensional Lévy processes*, arXiv preprint, 2011.
- [8] E. Lytvynov, *Orthogonal decompositions for Lévy processes with an application to the gamma, Pascal, and Meixner processes*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. 6 (2003), pp. 73–102.

- 
- [9] D. Nualart and W. Schoutens, *Chaotic and predictable representations for Lévy processes*, Stochastic Process. Appl. 90 (2000), pp. 109–122.
- [10] W. Schoutens, *Stochastic Processes and Orthogonal Polynomials*, Lecture Notes in Statist., Vol. 146, Springer, New York 2000.

Swansea University  
Department of Mathematics  
Singleton Park, Swansea, SA2 8PP, U.K.  
*E-mail:* 380535@swansea.ac.uk

Swansea University  
Department of Mathematics  
Singleton Park, Swansea, SA2 8PP, U.K.  
*E-mail:* e.lytvynov@swansea.ac.uk

*Received on 29.3.2013;*  
*revised version on 1.10.2013*

---