

Lazard's Theorem for Differential Algebraic Groups and Proalgebraic Groups

Marcin Chalupnik*
University of Warsaw

Piotr Kowalski †
University of Wrocław

Abstract

We prove that a differential group whose underlying variety is an affine space is unipotent. The problem is reduced to an infinite-dimensional version of Lazard's Theorem.

1 Introduction

It is well-known that a connected unipotent algebraic group is isomorphic as a variety to an affine space. Lazard ([La]) proved the converse: an algebraic group isomorphic as a variety to an affine space is unipotent. So, the algebraic structure of a unipotent algebraic group is determined by its geometry. Buium and Cassidy ([BC]) asked if the same holds for differential algebraic groups, i.e. groups defined by differential polynomials in some differential field. It was proved for groups of small dimension ($=1, 2$) by Cassidy ([Ca]), and for arbitrary groups over a differentially closed field of characteristic 0 by Kowalski and Pillay [KP]. The principal result of the present paper is the proof of this theorem in the full generality:

Theorem 1 *Suppose (K, D) is a differential field. Let G be a differential algebraic group over K , with underlying differential variety differentially isomorphic to A^n . Then G is unipotent (i.e. G may be embedded into a unipotent algebraic group).*

*Partially supported by Białynicki–Birula Subsidium of Foundation of Polish Science and the polish grant KBN 2 P03A 002 18.

†Partially supported by the polish grant KBN P03A 020 18.

The strategy of the proof of this theorem is borrowed from [KP]. We deduce Theorem 1 from a purely algebro-geometric, and interesting for its own right result, which was proved in [KP] for K an algebraically closed field of characteristic 0:

Theorem 2 *Let G be a group scheme over a field K whose underlying scheme is isomorphic to A^∞ (the projective limit of affine spaces). Then G is isomorphic in the category of group schemes over K to an inverse limit of unipotent algebraic groups.*

The paper is organized as follows. In Section 2 we reduce Theorem 1 to Theorem 2 (replacing model theory of [KP] by a simple argument using Hopf algebras). Theorem 1 is proved in Section 4. The proof utilizes in an essential way étale cohomology of schemes and we collect necessary facts concerning it in Section 3.

The reader may consult [BC] for background in differential algebraic geometry, while our main reference on étale cohomology is [Mi] (for basics see also very readable [Ta]).

The first author would like to thank Andrzej Weber for several enlightening comments concerning étale cohomology. The second author would like to thank Anand Pillay for suggesting this problem, and Ludomir Newelski for careful reading this paper and many helpful remarks.

2 The Reduction

Let us fix (K, D) a differential field. $K\{X\}$ denotes the ring of differential polynomials (in a set of variables X). As a ring $K\{X\} = K[X, DX, D^2X, \dots]$, where D^iX , $i \geq 0$, are tuples of new variables, and derivation on $K\{X\}$ extends $D^iX \mapsto D^{i+1}X$ ($D^0X := X$). A differential polynomial is an element of $K\{X\}$, and we naturally regard differential polynomials as functions from K^n into K . We recall some notions from differential algebraic geometry.

Definition 2.1

- i) A differential algebraic variety is a zero set of a finite number of differential polynomials.*
- ii) A morphism of differential algebraic varieties is a restriction of a differential polynomial function.*

iii) A differential algebraic group is a group object in the category of differential algebraic varieties.

Remark: Usually (e.g. [Bu], [BC]) a morphism of differential algebraic varieties is defined as a function locally given by differential rational functions rather than a globally defined polynomial. However on an affine space these two notions coincide, so we may adopt the more convenient definition.

Lemma 2.2 *If $f : G \longrightarrow H$ is a morphism between affine reduced group schemes, then there exist inverse systems (G_i) , (H_i) of algebraic groups and a morphism F between them, such that $G = \lim(G_i)$, $H = \lim(H_i)$ and $f = \lim(F)$.*

Proof: The proof becomes straightforward when we turn to the category of Hopf algebras, which is dual to the category of affine group schemes. The morphism f corresponds to a morphism $\phi : B \longrightarrow A$, where B is the Hopf algebra corresponding to H and A corresponds to G . Since any Hopf algebra is a direct limit of a system of its finitely generated Hopf subalgebras [Wa p. 24], there exist systems (A_i) , and (B_i) of finitely generated reduced Hopf algebras, such that $A = \text{colim}(A_i)$, and $B = \text{colim}(B_i)$. Note that finitely generated reduced Hopf algebras correspond to algebraic groups. Denote by ϕ_i^* the map ϕ restricted to the B_i . Then, since B_i is finitely generated, there exists n_i such that ϕ_i^* can be factorized through the map $\phi_i : B_i \longrightarrow A_{n_i}$. So $\phi = \lim(\phi_i)$, where $(\phi_i)_{i \in N}$ is a map between direct systems $(B_i)_{i \in N}$ and $(A_{n_i})_{i \in N}$. Going back to the category of affine group schemes we obtain the result. \square

Remark: A part of Lemma 2.2 (the existence of (G_i)) was proved in [KP] by a model-theoretical argument (valid in any stable theory after a suitable reformulation) for algebraic groups over an algebraically closed field.

Corollary 2.3 *Theorem 2 implies Theorem 1.*

Proof: Let us consider the ring $K\{X\}$ of differential polynomials in n variables. It is also a ring of differential regular functions (i.e. morphisms into A^1) on A^n , so as in algebraic geometry, differential group structure on A^n gives us the Hopf algebra structure on $K\{X\}$. We denote this Hopf algebra by A (it is also a differential algebra). It corresponds to an infinite-dimensional

group scheme G^* . The group $G^*(K)$ may be thought of as the set K^∞ with the group operation given by the sequence $(\mu, D(\mu), D^2(\mu), \dots)$, where D is a derivation in $K\{X\}$, and $K\{X\}$ acts on K^∞ as the ring of polynomials of infinitely many variables. Then the map $\phi : a \mapsto (a, Da, D^2a, \dots)$ is a homomorphism between G and $G^*(K)$. From Lemma 2.2, we have $G^* = \lim(G_i)$, where G_i 's are algebraic groups. Denote by H_i , the Hopf algebra of G_i . The Hopf algebra A is a direct limit of the system (H_i) and it is finitely generated as a differential algebra, so there exists n such that H_n differentially generates A . Then the composition morphism $G \rightarrow G_n$ is an embedding, since it induces epimorphism of algebras of differential regular functions. Using Theorem 1 and Lemma 2.2 we see that G_n is unipotent. \square

Remark:

- i) The construction of G^* is the same as in [KP]. However here we do not need our field to be differentially closed to check that we obtain a group.
- ii) If G is an algebraic group, then G^* coincides with the Buium's infinite prolongation, which is an inverse limit of so-called twisted jet spaces [Bu]. If D vanishes on the field of definition, then the infinite prolongation of an algebraic group is an inverse limit of the usual jet spaces.

3 Some Étale Cohomology of Group Schemes

The aim of this section is to prove some facts about étale cohomology of group schemes which will be needed in the proof of Theorem 2. Throughout this section \mathbf{k} is an algebraically closed field and all schemes are of finite type over \mathbf{k} . We are interested in étale cohomology groups with coefficients in the constant sheaf \mathbf{Z}/l where l is a prime distinct from the characteristic. We say that a scheme X is \mathbf{Z}/l -acyclic if it has cohomology of a point i.e. $H_{et}^i(X, \mathbf{Z}/l) = 0$ for $i > 0$, and $H_{et}^0(X, \mathbf{Z}/l) = \mathbf{Z}/l$; a morphism of schemes is called \mathbf{Z}/l -acyclic if it induces an isomorphism on $H_{et}^*(-, \mathbf{Z}/l)$. The main computational tool in étale cohomology theory is the Leray spectral sequence associated to a morphism of schemes $f : X \rightarrow Y$:

$$E_2^{ij} = H_{et}^i(X, \mathbf{R}^j f_*(\mathbf{Z}/l)) \Rightarrow H_{et}^{i+j}(X, \mathbf{Z}/l).$$

The typical application of the Leray spectral sequence is the following criterion of the acyclicity of a morphism:

Lemma 3.1 *Let $f : X \longrightarrow Y$ be either proper or smooth. Suppose all fibres of f are \mathbf{Z}/l -acyclic. Then f is \mathbf{Z}/l -acyclic.*

Proof: By the Proper (or Smooth) Base Change Theorem ([Mi] p. 224, p. 230) we may identify stalks of $\mathbf{R}^j f_*(\mathbf{Z}/l)$ with j -th cohomology of fibres. Thus all rows except 0-th in the Leray spectral sequence disappear. Now it suffices to observe that since f has connected fibre, then $\mathbf{R}^0 f_*(\mathbf{Z}/l) = \mathbf{Z}/l$. \square

If f is locally trivial (i.e. locally a projection from a product) and Y is smooth, then the Leray spectral sequence takes form known from algebraic topology. We recall that a scheme is simply connected if it has trivial algebraic fundamental group (i.e. the group classifying étale coverings of a scheme).

Lemma 3.2 *Suppose $f : X \longrightarrow Y$ is locally trivial in the étale topology, and Y is smooth, connected and simply connected. Denote by F the fiber of f . Then the Leray spectral sequence associated with f has the following form:*

$$E_2^{ij} = H_{et}^i(X, H_{et}^j(F, \mathbf{Z}/l)) \Rightarrow H_{et}^{i+j}(X, \mathbf{Z}/l),$$

where $H_{et}^j(F, \mathbf{Z}/l)$ is a constant sheaf.

Proof: From the Smooth Base Change Theorem ([Mi] p. 230) we derive that if f is a projection, then $\mathbf{R}^j f_*(\mathbf{Z}/l)$ may be identified with a constant sheaf having a stalk $H_{et}^j(F, \mathbf{Z}/l)$, where F is a fiber of f . So for f being locally trivial, the sheaf $\mathbf{R}^j f_*(\mathbf{Z}/l)$ is locally trivial with a stalk $H_{et}^j(F, \mathbf{Z}/l)$. However, analogously to the classical context, a locally constant sheaf with finite stalks (we point out that $H_{et}^j(F, \mathbf{Z}/l)$ for smooth F is always finite ([Mi] p. 244)) on a connected scheme Y is determined by an action of algebraic fundamental group of Y on a stalk ([Mi] p. 156). Thus for a simply connected scheme Y a locally constant sheaf with finite stalks must be constant and the Leray spectral sequence has the required form. \square

Another useful spectral sequence is the Hochschild–Serre spectral sequence corresponding to a Galois covering $p : X \longrightarrow Y$ with the structural group G ([Mi] p. 105):

$$E_2^{ij} = H^i(G, H^j(X, \mathbf{Z}/l)) \Rightarrow H^{i+j}(Y, \mathbf{Z}/l),$$

here $H^i(G, -)$ denotes cohomology of the discrete group G .
 Now we turn to the facts we need for the proof of Theorem 2.

Proposition 3.3 *Let $f : G \rightarrow H$ be an isogeny of connected affine algebraic groups. Then for almost all primes l , f is \mathbf{Z}/l -acyclic.*

Proof: Let K be the kernel of f regarded as a finite affine group scheme. Then it may be obtained as an extension: $1 \rightarrow K^0 \rightarrow K \rightarrow \Pi_0(K) \rightarrow 1$, where K^0 is connected and $\Pi_0(K)$ is étale ([Wa] p. 51). Hence we may factorize f as

$$G \xrightarrow{f_c} G/K^0 \xrightarrow{f_{et}} (G/K^0)/\Pi_0(K) = H,$$

where f_c has connected fibres and f_{et} is a Galois covering.

Since K^0 as a scheme is just a point (in general with multiplicities), fibres of f_c are \mathbf{Z}/l -acyclic. Moreover f_c being finite must be proper. Thus it satisfies the assumptions of Lemma 3.1, hence is \mathbf{Z}/l -acyclic for any l .

Let us turn to f_{et} and take l prime to $|\Pi_0(K)|$. Then the Hochschild–Serre spectral sequence corresponding to f_{et} collapses and we get $H_{et}^*(H, \mathbf{Z}/l) = (H_{et}^*((G/K^0), \mathbf{Z}/l))^{\Pi_0(K)}$, which gives us a monomorphism $H_{et}^*(H, \mathbf{Z}/l) \rightarrow H_{et}^*((G/K^0), \mathbf{Z}/l)$. This monomorphism is compatible with f_{et}^* by the very construction of the spectral sequence. Now it remains to show that $\Pi_0(K)$ acts trivially on $H_{et}^*((G/K^0), \mathbf{Z}/l)$. But the action of $\Pi_0(K)$ on G/K^0 extends to the action of the whole group G/K^0 , which is connected. Thus the triviality of the action at the level of cohomology will follow if we show that any embedding $\phi_g : G/K^0 \rightarrow G/K^0 \times G/K^0$ defined by $\phi_g(h) = (h, g)$ induces the same morphism on cohomology as ϕ_e . The last fact follows immediately from the Kunneth formula for $G/K^0 \times G/K^0$ ([SGA] p. 236), and the trivial observation that two constant morphisms into a connected scheme induce the same on étale cohomology. \square

Proposition 3.4 *Let G be a simple group. Then for almost all primes l , G is not \mathbf{Z}/l -acyclic.*

Proof: First observe that G has finite algebraic fundamental group. Indeed, it suffices to show that there is only a finite number of étale coverings of G . But this follows from the fact that any étale covering is étale (hence central) isogeny, while we have only a finite number of possibilities for centers of groups having root systems isomorphic to the root system of G ([Hu] p. 215), and all these centers are finite. Thus G has the universal covering being an

étale isogeny. Therefore, according to Proposition 3.3, we may assume that G is simply connected. Let $B \subset G$ be a Borel subgroup. Then the sequence $B \rightarrow G \rightarrow G/B$ is locally trivial in the étale topology [Se], and G/B is simply connected (this follows from $\Pi_1^{alg}(G) = 0$ and connectivity of B). Then, by Lemma 3.2 we have:

$$E_2^{ij} = H_{et}^i(G/B, H_{et}^j(B, \mathbf{Z}/l)) \Rightarrow H_{et}^{i+j}(G, \mathbf{Z}/l)$$

Now take $i_0 = \max\{i : H^i(G/B, \mathbf{Z}/l) \neq 0\}$. Since G/B is projective, it has the fundamental class ([Mi] pp. 247–252), so $i_0 = 2 \cdot \dim(G/B) > 0$. Similarly we take $j_0 = \max\{j : H^j(B, \mathbf{Z}/l) \neq 0\}$. Then we have $0 \neq E_2^{i_0 j_0} = E_\infty^{i_0 j_0}$, hence $H_{et}^{i_0+j_0}(G, \mathbf{Z}/l) \neq 0$, so G is not \mathbf{Z}/l -acyclic. \square

4 The proof of Theorem 2

The main ingredient in the proof is the following:

Proposition 4.1 *Let $f : G \rightarrow Q$ be an epimorphism of connected affine algebraic groups defined over an algebraically closed field \mathbf{k} , and assume that f factorizes through an affine space. Then Q is solvable.*

Proof: Suppose Q is not solvable. Then we have an epimorphism $p : Q \rightarrow S$ with simple S , hence we may assume that Q is already simple. According to the acyclicity of \mathbf{A}^n ([Mi] p. 295) and Proposition 3.4 in order to obtain contradiction it suffices to show that f induces a monomorphism on étale cohomology with coefficients in \mathbf{Z}/l for almost all l 's.

Let R_u be the unipotent radical of G . Then since $R_u \subset \ker(f)$, we may factorize f as $G \xrightarrow{g} G/R_u \xrightarrow{h} Q$. Let us first investigate g . Observe that g is smooth and has acyclic fibers, since R_u as a variety is isomorphic to an affine space. Thus according to Lemma 3.1 it induces an isomorphism on étale cohomology for any l prime to the characteristic. So, we may focus on h . Since G/R_u is reductive we have an isogeny $i : T \times S_1 \times \dots \times S_n \rightarrow G/R_u$, where T is a torus and each S_i is simple. We shall see that $(h \circ i)^*$ is a monomorphism on étale cohomology for some l . Indeed, since Q is simple, there exists such j that $(h \circ i)|_{S_j} : S_j \rightarrow Q$ is also an isogeny. Hence $((h \circ i)|_{S_j})^*$ is an isomorphism for almost all l 's, by Proposition 3.3. Thus $(h \circ i)^*$ (a fortiori h^*) is a monomorphism for these l 's. \square

In fact we need the following, quite straightforward, generalization of the above proposition:

Corollary 4.2 *Let K be any field, and $f : G \longrightarrow Q$ be an epimorphism of connected affine algebraic groups defined over K . Suppose that f factors through an affine space. Then Q is unipotent.*

Proof: Let $G_L = G \otimes L, Q_L = Q \otimes L, f_L = f \otimes L$, where L is the algebraic closure of K . Then the epimorphism $f_L : G_L \longrightarrow Q_L$ factors through an affine space, so by Lemma 4.1, Q_L is solvable. Moreover, any connected solvable group is a semidirect product of its unipotent radical R and some algebraic torus ([Hu] p. 123). But since Q_L is an image of an affine space, it cannot be mapped onto a torus. Thus Q_L is unipotent. It is also easy to see that Q_L is unipotent iff Q is (see e.g. [Wa] p. 64). This completes the proof. \square

Now we are in a position to prove our main theorem. We recall that by 2.5 it suffices to prove Theorem 2.

Proof of Theorem 2: Let G be a group scheme whose underlying scheme is A^∞ . By Lemma 2.2, G is isomorphic to an inverse limit of a system (G_i) of algebraic groups. It is enough to prove that G_i 's are unipotent. Fix a positive integer n . Then, like in the proof of Lemma 2.2, the isomorphism between G and $\lim(G_i)$ yields morphisms $f : G_N \longrightarrow A^l$, and $g : A^l \longrightarrow G_n$, for some l and $N > n$. The composition $g \circ f$ is an epimorphism, since it comes from the identity map on G . From Corollary 4.2, G_n is unipotent. \square

References

- [Bu] A. Buium, Geometry of differential polynomial functions I : Algebraic groups, *Amer. J. Math.* **115** (1993), 1385-1444.
- [BC] A. Buium, Ph.J. Cassidy, Differential algebraic geometry and differential algebraic groups, to appear in collected works of E. Kolchin.
- [Ca] Ph.J. Cassidy, Unipotent differential algebraic groups, in *Contributions to algebra*, Academic Press, 1977.

- [SGA] P. Deligne and J. F. Boutot, A. Grothendieck, L. Ilusie, J. L. Verdier, *SGA 4 $\frac{1}{2}$ Cohomologie Étale*, Lectures Notes in Math. 569, Springer, Heidelberg (1977).
- [Hu] J. Humphreys, *Linear Algebraic Groups*, Springer, New York (1981).
- [PK] P. Kowalski, A. Pillay, Proalgebraic and differential algebraic group structures on affine spaces, to appear in *Amer. J. Math.*
- [La] M. Lazard, Sur nilpotence de certains groupes algébriques, C.R.A.S. Paris **241** (1955), 1687-1689.
- [Mi] J. Milne, *Étale Cohomology*, Princeton University Press, Princeton (1980).
- [Se] J. P. Serre, Espaces fibrés algébriques, (Séminaire Chevalley 2 (1958), Exposé 1), Secrétariat Mathématique, Paris.
- [Ta] G. Tamme, *Introduction to Étale Cohomology*, Springer, Berlin (1994).
- [Wa] W. Waterhouse, *Introduction to Affine Group Schemes*, Springer, New York (1979).

INSTYTUT MATEMATYKI, UNIVERSITY OF WARSAW, UL. BANACHA 2,
02-097 WARSAW, POLAND
Electronic mail: MCHAL@MIMUW.EDU.PL

INSTYTUT MATEMATYCZNY, UNIVERSITY OF WROCLAW, PL. GRUN-
WALDZKI 2/4, 50-384 WROCLAW, POLAND
Electronic mail: PKOWA@MATH.UNI.WROC.PL