

A FORCING EXTENSION OF A SIMPLIFIED
 $(\omega_2, 1)$ MORASS WITH NO SIMPLIFIED $(\omega_2, 1)$
MORASS WITH LINEAR LIMITS

Franqui Cárdenas
Universidad Nacional de Colombia, Bogotá

July 14th, 2007
Wrocław

Old statement:using supercompact cardinals

Proof

New statement:using strongly unfoldable cardinals

Evidence

Proof

$\text{Con}(ZFC + \exists \kappa \text{ supercompact cardinal}) \implies$

$\text{Con}(ZFC + \exists(\omega_2, 1)\text{morass} + \neg\exists(\omega_2, 1) - \text{morass with linear limits})$

(Stanley)

Supercompact cardinals

Definition

κ is a θ -supercompact cardinal iff there exists $j : V \rightarrow M$ such that $cp(j) = \kappa$ and $M^\theta \subseteq M$.

κ is supercompact iff for all $\theta \in On$, κ is θ -supercompact.

Supercompactness $\implies V \neq L$.

Supercompact cardinals

Definition

κ is a θ -supercompact cardinal iff there exists $j : V \rightarrow M$ such that $cp(j) = \kappa$ and $M^\theta \subseteq M$.

κ is supercompact iff for all $\theta \in On$, κ is θ -supercompact.

Supercompactness $\implies V \neq L$.

Supercompact cardinals

Definition

κ is a θ -supercompact cardinal iff there exists $j : V \rightarrow M$ such that $cp(j) = \kappa$ and $M^\theta \subseteq M$.

κ is supercompact iff for all $\theta \in On$, κ is θ -supercompact.

Supercompactness $\implies V \neq L$.

Definition

κ regular cardinal. A simplified $(\kappa, 1)$ morass is a sequence $\langle \varphi_\xi; \mathcal{G}_{\xi\tau} : \xi < \tau \leq \kappa \rangle$ where

$$\mathcal{G}_{\xi\tau} = \{b : \varphi_\xi \rightarrow \varphi_\tau \mid b \text{ order preserving}\}$$

such that:

- ▶ $\varphi_\xi < \kappa$ and $|\mathcal{G}_{\xi\tau}| < \kappa$ for $\xi < \tau < \kappa$ and $\varphi_\kappa = \kappa^+$.
- ▶ Coherence.
- ▶ $\mathcal{G}_{\xi\xi+1} = \{id, f\}$ where f is a split function.
- ▶ If $\lim(\xi) \varphi_\xi = \bigcup_{\eta < \xi} \{b''\varphi_\eta \mid b \in \mathcal{G}_{\eta\xi}\}$.

Facts about morasses

- ▶ Simplified $(\kappa, 1)$ morasses implies the gap 2 cardinal theorem.
- ▶ There are simplified $(\omega, 1)$ morasses.
- ▶ If $V = L$ then for κ regular cardinal there are simplified $(\kappa, 1)$ morasses.
- ▶ Simplified $(\kappa, 1)$ morass implies $\square_{\kappa, \kappa}$.
- ▶ Simplified $(\kappa, 1)$ morass with linear limits implies \square_{κ} .

Facts about morasses

- ▶ Simplified $(\kappa, 1)$ morasses implies the gap 2 cardinal theorem.
- ▶ There are simplified $(\omega, 1)$ morasses.
- ▶ If $V = L$ then for κ regular cardinal there are simplified $(\kappa, 1)$ morasses.
- ▶ Simplified $(\kappa, 1)$ morass implies $\square_{\kappa, \kappa}$.
- ▶ Simplified $(\kappa, 1)$ morass with linear limits implies \square_{κ} .

Facts about morasses

- ▶ Simplified $(\kappa, 1)$ morasses implies the gap 2 cardinal theorem.
- ▶ There are simplified $(\omega, 1)$ morasses.
- ▶ If $V = L$ then for κ regular cardinal there are simplified $(\kappa, 1)$ morasses.
- ▶ Simplified $(\kappa, 1)$ morass implies $\square_{\kappa, \kappa}$.
- ▶ Simplified $(\kappa, 1)$ morass with linear limits implies \square_{κ} .

Facts about morasses

- ▶ Simplified $(\kappa, 1)$ morasses implies the gap 2 cardinal theorem.
- ▶ There are simplified $(\omega, 1)$ morasses.
- ▶ If $V = L$ then for κ regular cardinal there are simplified $(\kappa, 1)$ morasses.
- ▶ Simplified $(\kappa, 1)$ morass implies $\square_{\kappa, \kappa}$.
- ▶ Simplified $(\kappa, 1)$ morass with linear limits implies \square_{κ} .

Facts about morasses

- ▶ Simplified $(\kappa, 1)$ morasses implies the gap 2 cardinal theorem.
- ▶ There are simplified $(\omega, 1)$ morasses.
- ▶ If $V = L$ then for κ regular cardinal there are simplified $(\kappa, 1)$ morasses.
- ▶ Simplified $(\kappa, 1)$ morass implies $\square_{\kappa, \kappa}$.
- ▶ Simplified $(\kappa, 1)$ morass with linear limits implies \square_{κ} .

Proof using a supercompact cardinal

- ▶ Laver: If κ supercompact cardinal, then there is a forcing extension such that κ is still supercompact and it is indestructible under κ -directed closed forcings.
- ▶ The forcing which adds a simplified $(\kappa, 1)$ morass is κ -closed.
- ▶ Collapse κ to ω_2 .

Proof using a supercompact cardinal

- ▶ Laver: If κ supercompact cardinal, then there is a forcing extension such that κ is still supercompact and it is indestructible under κ -directed closed forcings.
- ▶ The forcing which adds a simplified $(\kappa, 1)$ morass is κ -closed.
- ▶ Collapse κ to ω_2 .

Proof using a supercompact cardinal

- ▶ Laver: If κ supercompact cardinal, then there is a forcing extension such that κ is still supercompact and it is indestructible under κ -directed closed forcings.
- ▶ The forcing which adds a simplified $(\kappa, 1)$ morass is κ -closed.
- ▶ Collapse κ to ω_2 .

$$V = L$$

- ▶ For any κ cardinal, \square_{κ} .
- ▶ For κ regular cardinal, there are $(\kappa, 1)$ -morasses (Jensen).
- ▶ Weakly compact cardinals, (strongly) unfoldable cardinals relativized to L .
- ▶ κ is weakly compact iff there is no $(\kappa, 1)$ -morass with linear limits (Donder).

$$V = L$$

- ▶ For any κ cardinal, \square_{κ} .
- ▶ For κ regular cardinal, there are $(\kappa, 1)$ -morasses (Jensen).
- ▶ Weakly compact cardinals, (strongly) unfoldable cardinals relativized to L .
- ▶ κ is weakly compact iff there is no $(\kappa, 1)$ -morass with linear limits (Donder).

$$V = L$$

- ▶ For any κ cardinal, \square_{κ} .
- ▶ For κ regular cardinal, there are $(\kappa, 1)$ -morasses (Jensen).
- ▶ Weakly compact cardinals, (strongly) unfoldable cardinals relativized to L .
- ▶ κ is weakly compact iff there is no $(\kappa, 1)$ -morass with linear limits (Donder).

$$V = L$$

- ▶ For any κ cardinal, \square_{κ} .
- ▶ For κ regular cardinal, there are $(\kappa, 1)$ -morasses (Jensen).
- ▶ Weakly compact cardinals, (strongly) unfoldable cardinals relativized to L .
- ▶ κ is weakly compact iff there is no $(\kappa, 1)$ -morass with linear limits (Donder).

Definition

Let κ be an inaccessible cardinal, M is a κ -model iff M is a transitive, $M \models ZF^-$, $|M| = \kappa$ with $\kappa \in M$ and $M^{<\kappa} \subseteq M$.

Definition

κ is θ -strongly unfoldable cardinal iff $\forall M (M \text{ } \kappa\text{-model} \implies \exists j, N [N \text{ transitive, } V_\theta \subseteq N, j : M \rightarrow N, cp(j) = \kappa, j(\kappa) \geq \theta])$.
(Villaveces)

Definition

Let κ be an inaccessible cardinal, M is a κ -model iff M is a transitive, $M \models ZF^-$, $|M| = \kappa$ with $\kappa \in M$ and $M^{<\kappa} \subseteq M$.

Definition

κ is θ -strongly unfoldable cardinal iff $\forall M (M \text{ } \kappa\text{-model} \implies \exists j, N [N \text{ transitive, } V_\theta \subseteq N, j : M \rightarrow N, cp(j) = \kappa, j(\kappa) \geq \theta])$.
(Villaveces)

Definition

κ is **strongly unfoldable** iff for all $\theta \in On$, κ is a θ -strongly unfoldable cardinal.

Fact: κ is weakly compact cardinal iff κ is κ -unfoldable cardinal.

Definition

κ is **strongly unfoldable** iff for all $\theta \in On$, κ is a θ -strongly unfoldable cardinal.

Fact: κ is weakly compact cardinal iff κ is κ -unfoldable cardinal.

Laver's preparation for other large cardinals

For κ strong, strongly compact, measurable and strongly unfoldable cardinals (Hamkins):

In all cases: **lottery preparation** relative to a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ is an ordinal arbitrary high below $j(\kappa)$.

Laver's preparation for other large cardinals

For κ strong, strongly compact, measurable and strongly unfoldable cardinals (Hamkins):

In all cases: **lottery preparation** relative to a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ is an ordinal arbitrary high below $j(\kappa)$.

Laver's preparation for other large cardinals

For κ strong, strongly compact, measurable and strongly unfoldable cardinals (Hamkins):

In all cases: **lottery preparation** relative to a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ is an ordinal arbitrary high below $j(\kappa)$.

If κ is strongly unfoldable cardinal, after the lottery preparation relative to f , κ strongly unfoldability is preserved by any $\mathbb{P} < \kappa$ -closed, κ -proper forcing (Hamkins, Johnstone)

$(2^{<\kappa} = \kappa)$ The forcing which adds a $(\kappa, 1)$ morass is κ -closed and κ^+ -c.c.

κ -properness forcing or preserving κ^+

If κ is strongly unfoldable cardinal, after the lottery preparation relative to f , κ strongly unfoldability is preserved by any $\mathbb{P} < \kappa$ -closed, κ -proper forcing (Hamkins, Johnstone)

($2^{<\kappa} = \kappa$) The forcing which adds a $(\kappa, 1)$ morass is κ -closed and κ^+ -c.c.

New statement using unfoldable cardinals

$\text{Con}(ZFC + \exists \kappa \text{ strongly unfoldable cardinal}) \implies$

$\text{Con}(ZFC + \exists(\omega_2, 1)\text{morass} + \neg\exists(\omega_2, 1) - \text{morass with linear limits})$

Proof:

- ▶ Let κ be strongly unfoldable cardinal and M a κ -model, there exists an embedding $j : M \rightarrow N$ with $cp(j) = \kappa$ and...
- ▶ Find a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ guess any value below $j(\kappa)$ (for free).
- ▶ Apply the lottery preparation to κ using f .
- ▶ Add the simplified $(\kappa, 1)$ morass. κ is still strongly unfoldable cardinal.
- ▶ Collapse κ to ω_2 .
- ▶ There is a simplified $(\omega_2, 1)$ morass but it is false \square_{ω_2} .

Proof:

- ▶ Let κ be strongly unfoldable cardinal and M a κ -model, there exists an embedding $j : M \rightarrow N$ with $cp(j) = \kappa$ and...
- ▶ Find a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ guess any value below $j(\kappa)$ (for free).
- ▶ Apply the lottery preparation to κ using f .
- ▶ Add the simplified $(\kappa, 1)$ morass. κ is still strongly unfoldable cardinal.
- ▶ Collapse κ to ω_2 .
- ▶ There is a simplified $(\omega_2, 1)$ morass but it is false \square_{ω_2} .

Proof:

- ▶ Let κ be strongly unfoldable cardinal and M a κ -model, there exists an embedding $j : M \rightarrow N$ with $cp(j) = \kappa$ and...
- ▶ Find a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ guess any value below $j(\kappa)$ (for free).
- ▶ Apply the lottery preparation to κ using f .
- ▶ Add the simplified $(\kappa, 1)$ morass. κ is still strongly unfoldable cardinal.
- ▶ Collapse κ to ω_2 .
- ▶ There is a simplified $(\omega_2, 1)$ morass but it is false \square_{ω_2} .

Proof:

- ▶ Let κ be strongly unfoldable cardinal and M a κ -model, there exists an embedding $j : M \rightarrow N$ with $cp(j) = \kappa$ and...
- ▶ Find a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ guess any value below $j(\kappa)$ (for free).
- ▶ Apply the lottery preparation to κ using f .
- ▶ Add the simplified $(\kappa, 1)$ morass. κ is still strongly unfoldable cardinal.
- ▶ Collapse κ to ω_2 .
- ▶ There is a simplified $(\omega_2, 1)$ morass but it is false \square_{ω_2} .

Proof:

- ▶ Let κ be strongly unfoldable cardinal and M a κ -model, there exists an embedding $j : M \rightarrow N$ with $cp(j) = \kappa$ and...
- ▶ Find a function $f : \kappa \rightarrow \kappa$ such that $j(f)(\kappa)$ guess any value below $j(\kappa)$ (for free).
- ▶ Apply the lottery preparation to κ using f .
- ▶ Add the simplified $(\kappa, 1)$ morass. κ is still strongly unfoldable cardinal.
- ▶ Collapse κ to ω_2 .
- ▶ There is a simplified $(\omega_2, 1)$ morass but it is false \square_{ω_2} .

Thanks!