

A technique for exponential change of measure for Markov processes.

Zbigniew Palmowski^{1,2,3}

Tomasz Rolski^{1,4}

September 9, 2002

Abstract

We consider a Markov process $X(t)$ with extended generator \mathbf{A} and domain $\mathcal{D}(\mathbf{A})$. Let $\{\mathcal{F}_t\}$ be a right-continuous history filtration and \mathbf{P}_t denote the restriction of \mathbf{P} to \mathcal{F}_t . Let $\tilde{\mathbf{P}}$ be another probability measure on (Ω, \mathcal{F}) such that $d\tilde{\mathbf{P}}_t/d\mathbf{P}_t = E^h(t)$, where

$$E^h(t) = \frac{h(X(t))}{h(X(0))} \exp \left(- \int_0^t \frac{(\mathbf{A}h)(X(s))}{h(X(s))} ds \right)$$

is a true martingale for a positive function $h \in \mathcal{D}(\mathbf{A})$. In this note, we demonstrate that the process $X(t)$ is a Markov process on the probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}, \tilde{\mathbf{P}})$, we find its extended generator $\tilde{\mathbf{A}}$ and provide sufficient conditions under which $\mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{D}(\mathbf{A})$. We apply this result to continuous time Markov chains, piecewise deterministic Markov processes and to diffusion processes (in this case a special choice of h yields the classical Cameron-Martin-Girsanov Theorem).

Keywords: Markov process, extended generator, exponential change of measure, Cameron-Martin-Girsanov theorem, (local) martingale, diffusion process, piecewise deterministic Markov process, Markov additive process.

AMS 1991 Subject Classification: Primary: 60G44, 60G25, 60J25, 60J35; Secondary: 60G48, 60H05, 60J57.

Bernoulli **8**, 2002.

¹Mathematical Institute, University of Wrocław, pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland

²EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

³This work was partially supported by KBN Grant No 5 P03 A 021 20 (2001-2003)

⁴This work was partially supported by KBN Grant No 2 P03 A 049 15 (1998-2001)

1 Introduction

The technique of exponential change of measure has been successfully applied in various theories such as large deviations (Schwartz and Weiss (1995), Ridder and Walrand (1992)), queues and fluid flows (Palmowski and Rolski (1996, 1998), Gautam *et al.* (1999), Kulkarni and Rolski (1994), Asmussen (1994,1995)), ruin theory (Asmussen (1994, 2000), Dassios and Embrecht (1989), Schmidli (1995, 1996, 1997a, 1997b)), simulation (Asmussen (1998), Ridder (1996)) or population genetics (Ethier and Kurtz (1993), Fukushima and Stroock (1986)). Typically, in the above books or papers, the process of interest is Markovian and, under a suitably chosen new probability measure, it is again a Markov process with some ‘nicer’ desired properties. To know parameters of the new process we need to find its generator. Although in some cases determining the new generator is straightforward, there are many situations which are more difficult. For such situations, a unified theory simplifies the calculations. In this note we present a detailed account of the change of probability measure technique, prepared for càdlàg Markov processes. We can also accommodate some non-Markovian processes that are Markovian with a supplementary component, e.g., piecewise deterministic Markov processes or Markov additive processes. For diffusion processes, a special case of the presented theory is a Girsanov type theorem.

Consider a Markov process $X(t)$ on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}, \mathbb{P})$ having extended generator \mathbf{A} with domain $\mathcal{D}(\mathbf{A})$. In this paper we use the martingale approach of Stroock and Varadhan (1979) to define generators. For each strictly positive function f , define

$$E^f(t) = \frac{f(X(t))}{f(X(0))} \exp \left(- \int_0^t \frac{(\mathbf{A}f)(X(s))}{f(X(s))} ds \right), \quad t \geq 0. \quad (1.1)$$

If for some function h , the process $E^h(t)$ is a martingale, then it is said to be an *exponential martingale*. In this case we call h a *good* function. Using this exponential martingale as a likelihood ratio process, we will define a new probability measure $\tilde{\mathbb{P}}$ on (Ω, \mathcal{F}) . In Theorem 4.2 we prove that under some mild assumptions, $X(t)$ is a Markov process on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}, \tilde{\mathbb{P}})$ having extended

generator

$$\tilde{\mathbf{A}}f = \frac{1}{h}[\mathbf{A}(fh) - f\mathbf{A}h] \quad (1.2)$$

and $\mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{D}(\mathbf{A})$. Note that (1.2) can be rewritten using the operator carré du champ (see e.g. Revuz and Yor (1991, Def. 3.2, p. 326)) $\tilde{\mathbf{A}}f = \mathbf{A}f + h^{-1}\langle h, f \rangle_{\mathbf{A}}$, where $\langle h, f \rangle_{\mathbf{A}} = \mathbf{A}(hf) - h\mathbf{A}f - f\mathbf{A}h$. Kunita (1969) and Ethier and Kurtz (1986) found that the two generators $\tilde{\mathbf{A}}$ and \mathbf{A} are related by $\tilde{\mathbf{A}} = \mathbf{A} + \mathbf{B}$, where \mathbf{B} is a linear operator. However, the domains in their papers are restricted to bounded functions. Under some additional assumptions, Ethier and Kurtz (1986, Th. 5.4, Th. 5.11 c) show that \mathbf{B} is the generator of some Markov process and they refer to $\tilde{\mathbf{A}}$ as a perturbation of \mathbf{A} . If a good function h is harmonic, that is $\mathbf{A}h = 0$ (or $\tilde{\mathbf{A}}h^{-1} = 0$), then $h(X(t))/h(X(0))$ is a martingale. In this case $\tilde{\mathbf{A}}f = h^{-1}\mathbf{A}(fh) - f\mathbf{A}h = h^{-1}\mathbf{A}(fh)$ (see Dynkin (1965), Chap. IX, (9.46) for a similar result expressed in terms of infinitesimal operators or Doob (1984), Chap. 2.VI.13 wherein the idea of h transforms was outlined).

In Section 5 we illustrate the theory applying it to several cases of special interest. For example, if $X(t)$ is a diffusion, then choosing the exponential function as a good function we obtain a Girsanov type theorem (see Theorem 5.5). We also show that a piecewise deterministic Markov process (PDMP), remains a PDMP under the new measure $\tilde{\mathbf{P}}$, and we find its characteristics (see Theorem 5.3). Other explicit forms of $\tilde{\mathbf{A}}$ are computed for continuous time Markov chains (CTMC) in Proposition 5.1, and for Markov additive processes (MAP) in Proposition 5.6.

Related material on change of measure or Girsanov type theorems can be found, for example, in Revuz and Yor (1991), Küchler and Sørensen (1997) and Jacod and Shiryaev (1987).

2 Preliminaries

2.1 Basic setup

Let E be a Borel space. Throughout the paper we denote the spaces of functions $f : E \rightarrow \mathbb{R}$ by: $\mathcal{M}(E)$ – the space of measurable functions and $\mathcal{C}(E)$ – the space of continuous functions. We add a subscript b to denote the restriction to bounded functions. We consider stochastic process

$X(t)$ assuming values in E and defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, equipped with a filtration $\{\mathcal{F}_t\}_{t \geq 0}$. We assume that $\mathcal{F} = \bigvee_{t \geq 0} \mathcal{F}_t$, i.e., \mathcal{F} is the smallest σ -field generated by all subsets of \mathcal{F}_t for all $t \geq 0$. We denote by \mathbb{F} the pair $(\mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0})$. We call (Ω, \mathbb{F}) a *filtered space* and $(\Omega, \mathbb{F}, \mathbb{P})$ a *filtered probability space*. For a probability measure \mathbb{P} we denote by \mathbb{P}_t its restriction to \mathcal{F}_t . All stochastic processes considered here are càdlàg.

Let $M(t)$ be a positive càdlàg martingale defined on a filtered probability space $(\Omega, \mathbb{F}, \mathbb{P})$ such that $\mathbb{E}M(0) = 1$. We define a family of measures $\tilde{\mathbb{P}}_t$ by

$$\frac{d\tilde{\mathbb{P}}_t}{d\mathbb{P}_t} = M(t) . \quad (2.3)$$

It can be proved that $\tilde{\mathbb{P}}_t$ is a consistent family, that is, for $s \leq t$ we have $\tilde{\mathbb{P}}_t(A) = \tilde{\mathbb{P}}_s(A)$ for all $A \in \mathcal{F}_s$.

We say that the *standard setup* is satisfied if

- (i) the filtered probability space $(\Omega, \mathbb{F}, \mathbb{P})$ has a right-continuous filtration $\{\mathcal{F}_t\}$,
- (ii) there exists a unique probability measure $\tilde{\mathbb{P}}$ such that $\tilde{\mathbb{P}}_t = \tilde{\mathbb{P}}|_{\mathcal{F}_t}$, where $\tilde{\mathbb{P}}_t$ is defined by (2.3).

The *standard setup* may not be satisfied if the filtration $\{\mathcal{F}_t\}$ is not separable (Parthasarathy (1967, Def. 2.1, Th. 4.2)); see e.g. the example of Rogers and Williams (1987, p. 83) and Karatzas and Shreve (1988, p. 193) concerning the augmented filtration of Brownian motion.

We now describe a *canonical model* for the *standard setup*. Assume that $\Omega \subset E^{[0, \infty)}$ is the space $D_E[0, \infty)$ of càdlàg functions from $[0, \infty)$ into E (in some cases we can also use $\mathcal{C}_E[0, \infty)$, the space of continuous functions into E) and consider the canonical process $X(t)$ on $(\Omega, \mathbb{F}, \mathbb{P})$ defined by $X(\omega, t) = \omega(t)$. The filtration is defined by $\{\mathcal{F}_t = \mathcal{F}_{t+}^X\}$, where $\mathcal{F}_t^X = \sigma\{X(s), s \leq t\}$ is the natural filtration generated by the process $X(t)$. We define a new family of probabilities via (2.3). By Kolmogorov theorem (see Stroock (1987, Th. 4.2, p. 106), Stroock and Varadhan (1997, Th. 1.3.5, p. 34), Parthasarathy (1967, Th. 4.2, p. 143)), there exists a unique probability measure $\tilde{\mathbb{P}}$ on (Ω, \mathbb{F}) such that $\tilde{\mathbb{P}}_t = \tilde{\mathbb{P}}|_{\mathcal{F}_t}$, where $\tilde{\mathbb{P}}_t$ fulfills (2.3). The above considerations are of course true

if we take the natural filtration $\{\mathcal{F}_t^X\}$ instead of $\{\mathcal{F}_{t+}^X\}$, but in applications often one needs the right-continuity assumption to define certain stopping times (e.g. moment of ruin in risk theory).

2.2 Markov processes

The following theorem was proved by Kunita and Watanabe (1963, Prop. 3 and 5). See also Dynkin (1965) and K  chler and S  rensen (1997, Prop. 6.3.5).

Theorem 2.1 *Let $X(t)$ be a Markov process on $(\Omega, \mathbb{F}, \mathbb{P})$. Assume that standard setup is satisfied. We define a new probability measure $\tilde{\mathbb{P}}$ by (2.3), where the martingale $M(t)$ is a nonnegative multiplicative functional for which $\mathbb{E}(M(0)) = 1$. Then on the new filtered probability space $(\Omega, \mathbb{F}, \tilde{\mathbb{P}})$, the process $X(t)$ is Markovian.*

For a Markov process $X(t)$ with values in E we define a *full (resp. extended) generator \mathbf{A}* by

$$\mathbf{A} = \{(f, f^*) \in \mathcal{M}(E) \times \mathcal{M}(E) : D^f(t) \text{ is a (resp. local) martingale}\} ,$$

where $D^f(t) = f(X(t)) - \int_0^t f^*(X(s)) ds$, $t \geq 0$ is *Dynkin's (resp. local) martingale* and the function $s \rightarrow f^*(X(s))$ is integrable \mathbb{P} -a.s. on $[0, t]$ for all $t \geq 0$. From now on we will identify all versions of the function f^* and we denote all these versions by $\mathbf{A}f$ if $(f, f^*) \in \mathbf{A}$.

By $\mathcal{D}(\mathbf{A})$ we denote the set of measurable functions $f \in \mathcal{M}(E)$ such that $\mathbf{A}f \in \mathcal{M}(E)$ exists, and such that $\int_0^t |\mathbf{A}f(X(s))| ds < \infty$ \mathbb{P} -a.s. for all $t \geq 0$. Thus

$$D^f(t) = f(X(t)) - \int_0^t \mathbf{A}f(X(s)) ds \tag{2.4}$$

is an \mathbb{F} -(local) martingale for $f \in \mathcal{D}(\mathbf{A})$. The space $\mathcal{D}(\mathbf{A})$ is called the *domain of the full (extended) generator \mathbf{A}* . If we restrict the domain of the extended or the full generator to a subset $\mathcal{D} \subset \mathcal{D}(\mathbf{A})$, then to avoid cumbersome notation we denote this subset also by $\mathcal{D}(\mathbf{A})$.

Remark 2.1 We will also use the notation $D^f(t)$ for the process given in (2.4) for any linear operator $\mathbf{A} : \mathcal{M}(E) \rightarrow \mathcal{M}(E)$, for which $s \rightarrow \mathbf{A}f(X(s))$ is integrable \mathbb{P} -a.s. on $[0, t]$ for all $t \geq 0$.

Remark 2.2 If $D^f(t)$ is a local martingale with respect to the augmented $\{\mathcal{F}_t^{\mathbf{P}}\}$ and $\{\mathcal{F}_t^{\mathbf{P}}\}$ -stopping times in a fundamental sequence are $\{\mathcal{F}_{t+}^X\}$ -stopping times, then $D^f(t)$ is also an $\{\mathcal{F}_{t+}^X\}$ -martingale. In this case, the extended generator and its domain do not change under augmentation.

Remark 2.3 Note that for $f \in \mathcal{D}(\mathbf{A})$ the process $\int_0^t \mathbf{A}f(X(s)) ds$ is a continuous process of finite variation. Thus the process $f(X(t))$ is a semimartingale. In particular, since the local martingale $D^f(t)$ is càdlàg this process is also càdlàg.

3 Exponential martingale

Consider a Markov process $X(t)$ defined on a filtered probability space $(\Omega, \mathbb{F}, \mathbb{P})$. For a linear operator $\mathbf{A} : \mathcal{M}(E) \rightarrow \mathcal{M}(E)$ we define

$$\mathcal{M}^*(\mathbf{A}) = \{f \in \mathcal{M}(E) : f(x) \neq 0 \text{ for all } x, \\ \int_0^t \left| \frac{(\mathbf{A}f)(X(s))}{f(X(s))} \right| ds < \infty \text{ and } \int_0^t |\mathbf{A}f(X(s))| ds < \infty \text{ for all } t \geq 0, \mathbb{P}\text{-a.s.}\}$$

Moreover, for an extended generator \mathbf{A} , we let

$$\mathcal{D}^*(\mathbf{A}) = \mathcal{M}^*(\mathbf{A}) \cap \mathcal{D}(\mathbf{A}) .$$

Thus, $E^f(t)$ defined in (1.1) makes sense for $f \in \mathcal{M}^*(\mathbf{A})$. The following result is slight extension of Proposition 3.2 in Ethier and Kurtz (1986, p. 175).

Lemma 3.1 *Let $f \in \mathcal{M}^*(\mathbf{A})$. Then the process $\{D^f(t), t \geq 0\}$ is a local martingale if and only if $\{E^f(t), t \geq 0\}$ is a local martingale.*

Proof. Assume that the process $D^f(t)$ is a local martingale. For $f \in \mathcal{M}^*(\mathbf{A})$, the process $\int_0^t \frac{\mathbf{A}f(X(s))}{f(X(s))} ds$ is continuous of finite variation. Hence by Jacod and Shiryaev (1987, Prop. 2.1.3, p. 75)

$$\left[f(X(t)), \exp \left(- \int_0^t \frac{(\mathbf{A}f)(X(s))}{f(X(s))} ds \right) \right]_t = 0 ,$$

where $[\cdot, \cdot]_t$ is a quadratic covariation. Using the integration-by-parts formula for semimartingales, we have

$$dE^f(t) = \frac{1}{f(X(0))} \exp\left(-\int_0^t \frac{(\mathbf{A}f)(X(s))}{f(X(s))} ds\right) dD^f(t) . \quad (3.5)$$

Thus, by Dellacherie and Meyer (1982, Th. VIII.3.e, p. 314), the process $E^f(t)$ is a local martingale. Now, assume that $E^f(t)$ is a local martingale. By (3.5) we also have that

$$dD^f(t) = f(X(0)) \exp\left(\int_0^t \frac{(\mathbf{A}f)(X(s))}{f(X(s))} ds\right) dE^f(t) ,$$

which completes the proof. □

Proposition 3.2 *Suppose that either one of the following two conditions holds for a positive function*

$h \in \mathcal{D}(\mathbf{A})$:

$$(M1) \quad h \in \mathcal{M}_b(E), \quad \text{and} \quad h^{-1} \mathbf{A}h \in \mathcal{M}_b(E) ,$$

or

$$(M2) \quad h, \mathbf{A}h \in \mathcal{M}_b(E) \quad \text{and} \quad \inf_x h(x) > 0 .$$

Then h is a good function.

Proof. A criterion of Protter (1990, Th. 47, p. 35), and Lemma 3.1 justify the above statement.

4 Main theorem

In this section we consider the Markov process from Section 3. Throughout this section, h is a good function and we define a new family of probabilities $\{\tilde{\mathbb{P}}_t, t \geq 0\}$ by

$$\frac{d\tilde{\mathbb{P}}_t}{d\mathbb{P}_t} = E^h(t), \quad t \geq 0 .$$

We now assume that the *standard setup* holds and thus there exists a unique probability measure $\tilde{\mathbb{P}}$ such that $\tilde{\mathbb{P}}_t = \tilde{\mathbb{P}}|_{\mathcal{F}_t}$. The Markovian property of $X(t)$ on the filtered probability space $(\Omega, \mathbf{F}, \tilde{\mathbb{P}})$ follows from Theorem 2.1.

We now quote the following useful lemma.

Lemma 4.1 [Liptser and Shiryaev (1986, Lem. 4.5.3, p. 188)] *An adapted process $Z(t)$ on a filtered space (Ω, \mathbf{F}) is a $\tilde{\mathbb{P}}$ -local martingale if $Z(t)E^h(t)$ is a \mathbb{P} -local martingale.*

For the operator $\tilde{\mathbf{A}}$ defined in (1.2), let

$$\mathcal{D}_{\mathbf{A}} = \left\{ f \in \mathcal{D}(\mathbf{A}) : hf \in \mathcal{D}^*(\mathbf{A}) \quad \text{and} \quad \int_0^t |\tilde{\mathbf{A}}f(X(s))| ds < \infty \text{ for all } t \geq 0, \tilde{\mathbb{P}}\text{-a.s.} \right\}.$$

For $f \in \mathcal{D}_{\mathbf{A}}$ we have $f(x) \neq 0$. Note also that if $f \in \mathcal{D}_{\mathbf{A}}$, then

$$\int_0^t \frac{(\tilde{\mathbf{A}}f)(X(s))}{f(X(s))} ds = \int_0^t \frac{(\mathbf{A}(fh))(X(s))}{fh(X(s))} ds - \int_0^t \frac{(\mathbf{A}h)(X(s))}{h(X(s))} ds$$

is well defined. Thus for $f \in \mathcal{D}_{\mathbf{A}}$ we can define the processes

$$\tilde{E}^f(t) = \frac{f(X(t))}{f(X(0))} \exp \left(- \int_0^t \frac{(\tilde{\mathbf{A}}f)(X(s))}{f(X(s))} ds \right) \quad (4.6)$$

and

$$\tilde{D}^f(t) = f(X(t)) - \int_0^t (\tilde{\mathbf{A}}f)(X(s)) ds.$$

We have

$$\begin{aligned} \tilde{E}^f(t)E^h(t) &= \frac{f(X(t))h(X(t))}{f(X(0))h(X(0))} \exp \left(- \int_0^t \left[\frac{(\tilde{\mathbf{A}}f)(X(s))}{f(X(s))} + \frac{(\mathbf{A}h)(X(s))}{h(X(s))} \right] ds \right) \\ &= \frac{fh(X(t))}{fh(X(0))} \exp \left(- \int_0^t \frac{(\mathbf{A}(fh))(X(s))}{fh(X(s))} ds \right). \end{aligned}$$

Applying Lemma 3.1 to $f \in \mathcal{D}_{\mathbf{A}}$ and Lemma 4.1 to $Z(t) = \tilde{E}^f(t)$, we have that $\tilde{D}^f(t)$ is a local martingale. Thus $\tilde{\mathbf{A}}$ is an extended generator of the process $X(t)$ on $(\Omega, \mathbf{F}, \tilde{\mathbb{P}})$ and $\mathcal{D}_{\mathbf{A}} \subset \mathcal{D}(\tilde{\mathbf{A}})$. Unfortunately, with this approach, we cannot easily identify the domain $\mathcal{D}(\tilde{\mathbf{A}})$ of the extended generator $\tilde{\mathbf{A}}$.

Let

$$\mathcal{D}_{\mathbf{A}}^h = \left\{ f \in \mathcal{D}(\mathbf{A}) : fh \in \mathcal{D}(\mathbf{A}) \quad \text{and} \quad \int_0^t |\tilde{\mathbf{A}}f(X(s))| ds < \infty \text{ for all } t \geq 0, \tilde{\mathbb{P}}\text{-a.s.} \right\}$$

and

$$\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \left\{ f \in \mathcal{D}(\tilde{\mathbf{A}}) : fh^{-1} \in \mathcal{D}(\tilde{\mathbf{A}}) \quad \text{and} \quad \int_0^t |\mathbf{A}f(X(s))| ds < \infty \text{ for all } t \geq 0, \mathbb{P}\text{-a.s.} \right\}.$$

Remark 4.1 Note that if $\inf_x h(x) > 0$ or $h \in \mathcal{C}(E)$, then by the inequality

$$|\tilde{\mathbf{A}}f| \leq \frac{|\mathbf{A}(fh)|}{|h|} + \frac{|\mathbf{A}h|}{|h|},$$

we have

$$\mathcal{D}_{\mathbf{A}}^h = \{f \in \mathcal{D}(\mathbf{A}) : fh \in \mathcal{D}(\mathbf{A})\}.$$

Similarly, if $h \in \mathcal{M}_b(E)$ or $h \in \mathcal{C}(E)$, then

$$\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \{f \in \mathcal{D}(\tilde{\mathbf{A}}) : fh^{-1} \in \mathcal{D}(\tilde{\mathbf{A}})\}.$$

Theorem 4.2 Suppose that $\mathcal{D}_{\mathbf{A}}^h = \mathcal{D}(\mathbf{A})$ and $\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \mathcal{D}(\tilde{\mathbf{A}})$. Then on the filtered probability space $(\Omega, \mathbb{F}, \tilde{\mathbb{P}})$ the process $X(t)$ is Markovian with extended generator $\tilde{\mathbf{A}}$ and $\mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{D}(\mathbf{A})$.

Proof. Let $f \in \mathcal{D}(\mathbf{A})$. Thus $f \in \mathcal{D}_{\mathbf{A}}^h$ and the process $\tilde{D}^f(t)$ is well defined. Moreover, it is càdlàg because by Remark 2.3, the process $f(X(t))$ is càdlàg. We will show that the process $\tilde{D}^f(t)$ is a $\tilde{\mathbb{P}}$ -local martingale. In that case we prove also inclusion $\mathcal{D}(\mathbf{A}) \subset \mathcal{D}(\tilde{\mathbf{A}})$. To prove the reverse inclusion, it suffices to change the measure by

$$\frac{d\mathbb{P}_t}{d\tilde{\mathbb{P}}_t} = E^h(t)^{-1},$$

which can be done since h is positive. Note that $h\tilde{\mathbf{A}}h^{-1} = -h^{-1}\mathbf{A}h$ and hence

$$\tilde{E}^{h^{-1}}(t) = \frac{h^{-1}(X(t))}{h^{-1}(X(0))} e^{-\int_0^t h\tilde{\mathbf{A}}h^{-1}(X(s)) ds} = (E^h(t))^{-1} \quad (4.7)$$

is a $\tilde{\mathbf{P}}$ -martingale. Note also that $f = 1 \in \mathcal{D}(\tilde{\mathbf{A}})$. So by the equality $\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \mathcal{D}(\tilde{\mathbf{A}})$, we have that $h^{-1} \in \mathcal{D}^*(\tilde{\mathbf{A}})$.

We prove now that the process $\tilde{D}^f(t)$ is a local martingale. Following Lemma 4.1, it suffices to show that the process $E^h(t)\tilde{D}^f(t)$ is a \mathbf{P} -local martingale. By the integration-by-parts formula for semimartingales, we get

$$\tilde{D}^f(t)E^h(t) = \int_0^t \tilde{D}^f(s-) dE^h(s) + \int_0^t E^h(s-) d\tilde{D}^f(s) + [E^h, \tilde{D}^f]_t .$$

However, from the definition of the process $\tilde{D}^f(t)$ and the operator $\tilde{\mathbf{A}}$, we have

$$\tilde{D}^f(t) = D^f(t) - \int_0^t \frac{\langle f, h \rangle_{\mathbf{A}}(X(s))}{h(X(s))} ds .$$

Hence

$$\begin{aligned} \tilde{D}^f(t)E^h(t) &= \int_0^t \tilde{D}^f(s-) dE^h(s) + \int_0^t E^h(s-) dD^f(s) \\ &\quad + [E^h, \tilde{D}^f]_t - \int_0^t E^h(s-) \frac{\langle f, h \rangle_{\mathbf{A}}(X(s))}{h(X(s))} ds . \end{aligned} \quad (4.8)$$

The first two components are local martingales. We now consider the third component. Let X^c be the continuous component, and X^d be the pure-jump component of the process $X(t)$, and let $\Delta f(X(t)) = f(X(t)) - f(X(t-))$. Then we have

$$d[E^h, \tilde{D}^f]_t = d[f(X), E^h]_t - d\left[\int_0^t \tilde{\mathbf{A}}f(X(s)) ds, E^h(t)\right]_t .$$

Note that $\int_0^t \tilde{\mathbf{A}}f(X(s)) ds$ is continuous of finite variation and hence $\left[\int_0^t \tilde{\mathbf{A}}f(X(s)) ds, E^h(t)\right]_t = 0$. Thus $d[E^h, \tilde{D}^f]_t = d[f(X), E^h]_t$, which by (3.5), is equal to

$$d\left[f(X(t)), \int_0^t \frac{E^h(s-)}{h(X(s-))} dh(X(s))\right]_t - d\left[f(X(t)), \int_0^t \frac{E^h(s-)}{h(X(s-))} \mathbf{A}h(X(s)) ds\right]_t . \quad (4.9)$$

Note that

$$\int_0^t \left| \frac{1}{h(X(s-))} E^h(s-) \mathbf{A}h(X(s)) \right| ds \leq \frac{1}{h(X(0))} \exp\left\{ \int_0^t \left| \frac{\mathbf{A}h(X(s))}{h(X(s))} \right| ds \right\} \int_0^t |\mathbf{A}h(X(s))| ds ,$$

which is finite \mathbb{P} -a.s. for all $t \geq 0$. Thus process $\int_0^t \frac{1}{h(X(s-))} E^h(s-) \mathbf{A} h(X(s)) ds$ is continuous of finite variation and the second component of (4.9) is equal to 0. We have

$$\begin{aligned} d[E^h, \tilde{D}^f]_t &= d \left[f(X^c(t)) + f(X^d(t)), \int_0^t \frac{1}{h(X(s-))} E^h(s-) d(h(X^c(s)) + h(X^d(s))) \right]_t \\ &= \frac{1}{h(X(t-))} E^h(t-) d[f(X^d(t)), h(X^d(t))]_t = \frac{1}{h(X(t-))} E^h(t-) d[f(X(t)), h(X(t))]_t . \end{aligned}$$

Using the integration-by-parts formula for semimartingales again, one obtains

$$\begin{aligned} d[E^h, \tilde{D}^f]_t &= \frac{1}{h(X(t-))} E^h(t-) d \left(f(X(t)) h(X(t)) - \int_0^t f(X(s-)) dh(X(s)) \right. \\ &\quad \left. - \int_0^t h(X(s-)) df(X(s)) \right) = \frac{1}{h(X(t-))} E^h(t-) dD^{fh}(t) \\ &\quad - \frac{1}{h(X(t-))} E^h(t-) f(X(t-)) dD^h(t) - \frac{1}{h(X(t-))} E^h(t-) h(X(t-)) dD^f(t) \\ &\quad + \frac{1}{h(X(t-))} E^h(t-) \left((\mathbf{A}(fh))(X(t)) - f(X(t-))(\mathbf{A}h)(X(t)) - h(X(t-))(\mathbf{A}f)(X(t)) \right) dt . \end{aligned}$$

The first three components above are all local martingales since they are the integrals with respect to local martingales. The last component is a Riemann differential, and is equal to

$$\frac{\langle f, h \rangle_{\mathbf{A}}(X(t))}{h(X(t))} E^h(t-) dt ,$$

which completes the proof in view of (4.8). □

5 Examples

5.1 Continuous time Markov chain

The simplest case we can analyze is when $X(t)$ is a CTMC. We will only consider CTMCs with a finite state space. The modification to a countable state space is straightforward, but it would require many additional assumptions.

Let $\mathbf{Q} = (q_{ij})_{i,j=1,\dots,\ell}$ be the intensity matrix of the process, which, in the case of the finite state space, is also the generator of the process. Here, functions f, h are column vectors. We define

$fh = (f_1 h_1, \dots, f_\ell h_\ell)^T$. Let h be positive. Now we have the following proposition (see also Rolski *et al.* (1999, Lem. 12.3.3)).

Proposition 5.1 *The new generator is $\tilde{Q} = (\tilde{q}_{ij})$, where*

$$\tilde{q}_{ij} = \begin{cases} q_{ij} \frac{h_j}{h_i}, & i \neq j \\ -\sum_{k \neq i} q_{ik} \frac{h_k}{h_i}, & i = j \end{cases}.$$

Proof. By inspection.

Corollary 5.2 *If Q is irreducible and reversible with the stationary probability vector π , then the stationary vector $\tilde{\pi}$ for \tilde{Q} is given by*

$$\tilde{\pi}_i = C \frac{\pi_i h_i}{\prod_{j \neq i} h_j},$$

where C is a constant such that $\sum_{j=1}^\ell \tilde{\pi}_j = 1$.

5.2 Piecewise Deterministic Markov Process

We follow Davis' (1993) description of PDMPs. Let E be a state space consisting of pairs $x = (\nu, z)$, where ν assumes a finite number of values from a set \mathcal{I} and z belongs to an open subset \mathcal{O}_ν of $\mathbb{R}^{d(\nu)}$ (note that E is a Borel space). The process $X(t)$ is determined (see Davis (1993, p. 62)), by the following:

1. $\mathcal{X} = \sum_{i=1}^{d_\nu} g^{\nu,i}(x) \frac{\partial}{\partial z_i}$, a vector field determining the flow $\phi_\nu(t, z)$ (we assume that $g^{\nu,i}$ are locally Lipschitz continuous),
2. $\lambda(\cdot)$, a force of transitions,
3. $Q(\cdot, \cdot)$, a transition kernel.

Denote by $\partial \mathcal{O}_\nu$ the boundary of \mathcal{O}_ν and let

$$\partial^* \mathcal{O}_\nu = \{z \in \partial \mathcal{O}_\nu : z = \phi_\nu(t, z') \text{ for some } (t, z') \in \mathbb{R}_+ \times \mathcal{O}_\nu\},$$

$$\Gamma = \{(\nu, z) \in \partial E : \nu \in \mathcal{I}, z \in \partial^* \mathcal{O}_\nu\},$$

$$t^*(\nu, z) = \sup\{t > 0 : \phi_\nu(t, z) \text{ exists and } \phi(t, z) \in \mathcal{O}_\nu\} .$$

The set Γ is called an *active boundary*. It is the set of boundary points of E , which can be reached from E via integral curves in finite time. For each point (ν, z) , $t^*(\nu, z)$ is the time needed to reach the boundary from (ν, z) . We will assume that $\phi_\nu(t^*(\nu, z), z) \in \Gamma$ if $t^*(\nu, z) < \infty$, which means that the integral curves cannot "disappear" inside E . Assume that

$$\lim_{n \rightarrow \infty} T_n = \infty, \quad \mathbb{P}\text{-a.s.} \quad (5.10)$$

where T_n denotes the consecutive jumps of the process $X(t)$. To get (5.10) we can assume that λ is bounded and that one of the following conditions is satisfied: $t^*(x) = \infty$ for each $x = (\nu, z) \in E$, that is, there are no active boundary points (e.g. $\Gamma = \emptyset$), or for some $\epsilon > 0$, we have $Q(x, B_\epsilon) = 1$ for all $x \in \Gamma$, where $B_\epsilon = \{x \in E : t^*(x) \geq \epsilon\}$. The last condition means that the minimal distance between consecutive boundary hitting times is not smaller than ϵ (see Davis (1993, Prop. 24.6, p. 60)).

From Davis (1993, Prop. 26.14, p. 69), the formula for the extended generator is

$$\mathbf{A}f(x) = \mathcal{X}f(x) + \lambda(x) \int_E (f(y) - f(x))Q(x, dy) ,$$

where $x \in E$ and the domain $\mathcal{D}(\mathbf{A})$ consists of every function f that is the restriction to E of a measurable function $\bar{f} : E \cup \Gamma \rightarrow \mathbb{R}$ satisfying three conditions:

(i) for each $(\nu, z) \in E$ the function $t \rightarrow \bar{f}(\nu, \phi_\nu(t, z))$ is absolutely continuous on $(0, t^*(\nu, z))$,

(ii) for each $x \in \Gamma$,

$$\bar{f}(x) = \int_E \bar{f}(y)Q(x, dy) , \quad (5.11)$$

(iii) for $n = 1, 2, \dots$,

$$\mathbb{E} \left(\sum_{i=1}^n |\bar{f}(X(T_i)) - \bar{f}(X(T_i -))| \right) < \infty . \quad (5.12)$$

From (1.2) we have

$$(\tilde{\mathbf{A}}f)(x) = (\mathcal{X}f)(x) + \frac{\lambda(x)H(x)}{h(x)} \int_E (f(y) - f(x)) \frac{h(y)}{H(x)} Q(x, dy) ,$$

where $H(x) = \int_E h(y)Q(x, dy)$. We assume that

$$\lim_{n \rightarrow \infty} T_n = \infty, \quad \tilde{\mathbb{P}}\text{-a.s.} \quad (5.13)$$

This condition holds if, for example, $\lambda(x)$ is bounded, $\inf_x h(x) > 0$ and $\Gamma = \emptyset$. We can now conclude.

Theorem 5.3 *Assume that h is a good function satisfying $H(x) < \infty$ for all $x \in E$. Suppose that (5.13) holds, and that $\mathcal{D}_{\mathbf{A}}^h = \mathcal{D}(\mathbf{A})$ and $\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \mathcal{D}(\tilde{\mathbf{A}})$. Then on the new probability space $(\mathcal{D}_E[0, \infty), \mathbb{F}, \tilde{\mathbb{P}})$, the process $X(t)$ is a PDMP with the unchanged differential operator \mathcal{X} and the following jump intensity and transition kernel:*

$$\tilde{\lambda}(x) = \frac{\lambda(x)H(x)}{h(x)}, \quad \tilde{Q}(x, dy) = \frac{h(y)}{H(x)}Q(x, dy). \quad (5.14)$$

The domain does not change: $\mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{D}(\mathbf{A})$.

Remark 5.1 Assume that condition (M1) or (M2) of Proposition 3.2 holds. Then h is a good function and $H(x) < \infty$ for all x . Moreover if, for each $f \in \mathcal{D}(\mathbf{A})$ the function fh satisfies condition (5.11) (e.g. if there are no active boundary points: $\Gamma = \emptyset$), then $\mathcal{D}_{\mathbf{A}}^h = \mathcal{D}(\mathbf{A})$ and $\mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \mathcal{D}(\tilde{\mathbf{A}})$ by Remark 4.1. Thus, if (5.13) holds additionally, then all assumptions of Theorem 5.3 are satisfied. In particular, this is the case when: $\Gamma = \emptyset$, $\lambda(x)$ is bounded and h is an absolutely continuous, bounded function such that $\inf_x h(x) > 0$.

Remark 5.2 Consider a positive function $h \in \mathcal{D}(\mathbf{A})$ which is harmonic (typically unbounded so that conditions (M1) and (M2) of Proposition 3.2 do not hold) and which belongs to the domain of the full generator of the process $X(t)$. That is, h is a good function and by Davis (1993, Rem. 26.17, p. 70), it satisfies a condition stronger than (5.12), namely

$$\mathbb{E} \left(\sum_{T_n \leq t} |\bar{h}(X(T_i)) - \bar{h}(X(T_i-))| \right) < \infty, \quad (5.15)$$

for all $t \geq 0$. Assume that (5.13) holds, $H(x) < \infty$ for all $x \in E$ and fh satisfies (5.11) (e.g. $\Gamma = \emptyset$). We restrict the domains of the extended generators \mathbf{A} and $\tilde{\mathbf{A}}$ to the bounded measurable functions satisfying (i) and (ii) and we denote these sets also by $\mathcal{D}(\mathbf{A})$ and $\mathcal{D}(\tilde{\mathbf{A}})$, respectively. Then by Theorem 5.3, the process $X(t)$ on $(\mathcal{D}_E[0, \infty), \mathbb{F}, \tilde{\mathbb{P}})$ is a PDMP with the parameters given in (5.14) and $\mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{D}(\mathbf{A})$. Such a framework is often used in risk theory (see e.g. Rolski *et al.* (1999, p. 460)).

5.3 Diffusion processes

Let $\mathbf{a} = (a_{ij})_{i,j=1,\dots,d}$ and $b = (b_1, \dots, b_d)^T$ be functions of $x \in \mathbb{R}^d$ satisfying

1. $b_i \in \mathcal{C}(\mathbb{R}^d)$ and for some $L > 0$,

$$|b_i(x)| \leq L(1 + \|x\|), \quad x \in \mathbb{R}^d, \quad (5.16)$$

2. $\mathbf{a}(x)$ is a strictly positive definite matrix and

$$a_{ij} \in \mathcal{C}_b(\mathbb{R}^d). \quad (5.17)$$

In this section we assume that $X(t)$ is a Markov diffusion process on the state space $E = \mathbb{R}^d$ with a given initial state $X(0)$. We consider here $\Omega = \mathcal{C}_{\mathbb{R}^d}[0, \infty)$ and $X(\omega, t) = \omega(t)$ with the right-continuous filtration $\{\mathcal{F}_t = \mathcal{F}_{t+}^X\}_{t \geq 0}$. The extended generator \mathbf{A} of the diffusion process is

$$(\mathbf{A}f)(x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2 f(x)}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(x) \frac{\partial f(x)}{\partial x_i}, \quad (5.18)$$

where $x \in \mathbb{R}^d$ and the family of twice continuously differentiable functions $f \in \mathcal{C}^2(\mathbb{R}^d)$ are included in the domain of this generator $\mathcal{D}(\mathbf{A})$; see Karatzas and Shreve (1988, Prop. 4.2, p. 312), Rogers and Williams (1987, Th. 24.1, p. 170). We will write $\mathcal{D}(\mathbf{A}) = \mathcal{C}^2(\mathbb{R}^d)$. Let $c(x) = (c_1(x_1), \dots, c_d(x_d))^T : \mathbb{R}^d \rightarrow \mathbb{R}^d$, where

$$c_i \in \mathcal{C}_b^1(\mathbb{R}), \quad i = 1, \dots, d. \quad (5.19)$$

For $x_0 = (x_{0,1}, \dots, x_{0,d})$, $x = (x_1, \dots, x_d)$ and $y = (y_1, \dots, y_d)$ we define

$$\int_{x_0}^x c(y) dy = \sum_{j=1}^d \int_{x_{0,j}}^{x_j} c_j(y_j) dy_j .$$

We use the following function $h : \mathbb{R}^d \rightarrow \mathbb{R}_+$:

$$h(x) = \exp \left(\int_{x_0}^x c(y) dy \right) . \quad (5.20)$$

For $d = 1$, we have $h(x) = \exp(\int_{x_0}^x c(y) dy)$. Note that the function $h(x)$ satisfies:

$$\frac{1}{h(x)} \frac{\partial h(x)}{\partial x_j} = c_j(x_j) . \quad (5.21)$$

Moreover, we have that $h \in \mathcal{D}(\mathbf{A})$ and $h(x) \neq 0$. Hence $h \in \mathcal{D}^*(\mathbf{A})$ and thus by Lemma 3.1, the process $E^h(t)$ is a local martingale. However, a stronger result is also true. We adopt the convention that

$$(c^T b)(x) = c(x)^T b(x), \quad (\mathbf{a}c)(x) = \mathbf{a}(x)c(x), \quad (c^T \mathbf{a}c)(x) = c(x)^T \mathbf{a}(x)c(x) .$$

Lemma 5.4 *We have*

$$E^h(t) = \exp \left(\int_0^t c(X(s)) dX(s) - \int_0^t (c^T b)(X(s)) ds - \frac{1}{2} \int_0^t (c^T \mathbf{a}c)(X(s)) ds \right) , \quad (5.22)$$

where

$$\int_0^t c(X(s)) dX(s) = \sum_{j=1}^d \int_0^t c_j(X_j(s)) dX_j(s) .$$

Proof. From (5.18) and (5.21) we have

$$\frac{\mathbf{A}h(x)}{h(x)} = \frac{1}{2} \sum_{i=1}^d a_{ii}(x) c'_i(x_i) + \frac{1}{2} (c^T \mathbf{a}c)(x) + (c^T b)(x) , \quad (5.23)$$

where $c'_i(t) = \frac{d}{dz} c_i(z)|_{z=t}$. Then

$$\begin{aligned} E^h(t) &= \exp \left(\int_{X(0)}^{X(t)} c(x) dx \right) \exp \left(-\frac{1}{2} \int_0^t \sum_{i=1}^d a_{ii}(X(s)) c'_i(X_i(s)) ds \right) \\ &\quad \exp \left(-\frac{1}{2} \int_0^t (c^T \mathbf{a}c)(X(s)) dX(s) - \int_0^t (c^T b)(X(s)) ds \right) . \end{aligned}$$

Thus it suffices to show that

$$\sum_{i=1}^d \int_{X_i(0)}^{X_i(t)} c_i(z) dz = \sum_{i=1}^d \int_0^t c_i(X_i(s)) dX_i(s) + \frac{1}{2} \int_0^t \sum_{i=1}^d a_{ii}(X(s)) c'_i(X_i(s)) ds . \quad (5.24)$$

From Stroock (1987, p. 75), we have

$$d\langle X_i(t) - \int_0^t b_i(X(s)) ds \rangle_t = a_{ii}(X(t)) dt .$$

Now (5.24) follows from Ito's formula applied to the function $g(x) = \int_{x_0}^x c_i(z) dz$:

$$\begin{aligned} \int_{X_i(0)}^{X_i(t)} c_i(z) dz &= \int_0^t c_i(X_i(s)) dX_i(s) + \frac{1}{2} \int_0^t \frac{d}{dx} c'_i(X_i(s)) d\langle X_i(s) \rangle_s \\ &= \int_0^t c_i(X_i(s)) dX_i(s) + \frac{1}{2} \int_0^t c'_i(X_i(s)) d\langle X_i(s) - \int_0^s b_i(X(v)) dv \rangle_s \\ &= \int_0^t c_i(X_i(s)) dX_i(s) + \frac{1}{2} \int_0^t a_{ii}(X(s)) c'_i(X_i(s)) ds , \end{aligned} \quad (5.25)$$

where (5.25) follows from the fact that $\int_0^t b_i(X(s)) ds$ has the finite variation ($b_i \in \mathcal{C}(\mathbb{R}^d)$).

□

Note that by (5.17), (5.19) and Stroock (1987, p. 75) we have

$$\langle \int_0^t c(X(s)) dX(s) - \int_0^t c(X(s)) b(X(s)) ds \rangle_t = \int_0^t (c^T \mathbf{a} c)(X(s)) ds \leq K_t , \quad (5.26)$$

for some constant K_t . Thus from Rogers and Williams (1987, Th. 37.8, p. 77), the process $E^h(t)$ is a true martingale, that is, h is a good function.

As an illustration of the derived Theorem 4.2, we give another proof of the Cameron-Martin-Girsanov Theorem. In this theorem, we consider the new probability measure $\tilde{\mathbb{P}}$ defined by the martingale $d\tilde{\mathbb{P}}_t/d\mathbb{P}_t = E^h(t)$, where h is a good function of the form (5.20). We have

$$(\tilde{\mathbf{A}}f)(x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2 f(x)}{\partial x_i \partial x_j} + \sum_{i=1}^d \left(b_i(x) + \sum_{j=1}^d a_{ij}(x) \frac{1}{h(x)} \frac{\partial h(x)}{\partial x_j} \right) \frac{\partial f(x)}{\partial x_i} . \quad (5.27)$$

Theorem 5.5 *Suppose that h is a good function of the form (5.20). On the new probability space $(\mathcal{C}_{\mathbb{R}^d}[0, \infty), \mathbb{F}, \tilde{\mathbb{P}})$, the process $X(t)$ is a diffusion process with parameters*

$$\tilde{\mathbf{a}} = \mathbf{a}, \quad \tilde{\mathbf{b}} = \mathbf{b} + \mathbf{a} c .$$

Proof. The assumptions of Theorem 4.2 are satisfied, that is, $\mathcal{D}_{\mathbf{A}}^h = \mathcal{D}(\mathbf{A}) = \mathcal{D}_{\tilde{\mathbf{A}}}^{h^{-1}} = \mathcal{D}(\tilde{\mathbf{A}}) = \mathcal{C}^2(\mathbb{R}^d)$ and we deduce $\tilde{\mathbf{a}}$ and \tilde{b} from the form of the generator given in (5.27). □

Example 5.1 Take $d = 1$ and $c_i(x) = -\alpha$. In this case a diffusion process with infinitesimal variance function $a(x)$ and infinitesimal drift function $b(x)$ has new parameters $\tilde{a}(x) = a(x)$ and $\tilde{b}(x) = b(x) - \alpha a(x)$ after the exponential change of measure. In particular, if we consider Brownian motion $B(t)$, then after the exponential change of measure, the process $B(t)$ is Brownian motion with drift $-\alpha$ (see Itô and Watanabe (1965)).

Example 5.2 The second example concerns the so-called Gauss-Markov process $X(t)$ (see Karatzas and Shreve (1988, p. 355), which is the solution of the SDE:

$$dX(t) = \mathbf{m}X(t)dt + \boldsymbol{\sigma}dB(t) ,$$

where $\mathbf{m}, \boldsymbol{\sigma}$ are $d \times d$ matrices and $B(t)$ is d -dimensional Brownian motion. We assume that all eigenvalues of \mathbf{m} have strictly negative real parts. Then by Stroock (1987, Th. 2.6, p. 91) the process $X(t)$ is a diffusion with

$$\mathbf{a}(x) = \boldsymbol{\sigma}\boldsymbol{\sigma}^T, \quad b(x) = \mathbf{m}x ,$$

where $x = (x_1, \dots, x_d)^T$. After the exponential change of measure, the process $X(t)$ is a diffusion with parameters $\tilde{\mathbf{a}}(x) = \mathbf{a}(x)$ and $\tilde{b}(x) = \mathbf{m}x + \mathbf{a}c(x)$, where $c(x) = (c_1(x_1), \dots, c_d(x_d))^T$ and $c_i(x_i) \in \mathcal{C}_b^1(\mathbb{R})$. In particular, when $d = 1$, $\mathbf{m} = -\alpha \in \mathbb{R}$ and $\boldsymbol{\sigma} = \sqrt{a} \in \mathbb{R}_+$, then the Gauss-Markov process is a one-dimensional Ornstein-Uhlenbeck process. Taking $c(x) = c \in \mathbb{R}$, after the exponential change of measure, the process $X(t)$ is the diffusion with infinitesimal drift changed from $-\alpha x$ to $-\alpha x + ac$; see Kulkarni and Rolski (1994).

5.4 Markov Additive Process

Following Asmussen and Kella (2000) we consider a Markov additive process (MAP) $X(t)$, where $X(t) = X^{(1)}(t) + X^{(2)}(t)$, and the independent processes $X^{(1)}(t)$ and $X^{(2)}(t)$ are specified by the characteristics: $q_{ij}, G_i, \sigma_i, a_i, \nu_i(dx)$ which will be defined below. Let $J(t)$ be a right continuous irreducible, finite state space CTMC, with $\mathcal{I} = \{1, \dots, N\}$, and with the intensity matrix $\mathbf{Q} = (q_{ij})$. We denote the jumps of the process $J(t)$ by $\{T_i\}$ (with $T_0 = 0$). Let $\{U_n^i\}$ be i.i.d. random variables with distribution function $G_i(\cdot)$. Define the jump process by

$$X^{(1)}(t) = \sum_{n \geq 1} \sum_i U_n^i \mathbb{1}_{\{J(T_{n-1})=i, T_n \leq t\}}.$$

For each $i \in \mathcal{I}$, let $X^i(t)$ be a Lévy process such that

$$\log \mathbb{E} e^{\alpha X^i(1)} = \psi_i(\alpha) = a_i \alpha + \frac{\sigma_i^2 \alpha^2}{2} + \int_{-\infty}^{\infty} (e^{\alpha y} - 1 - \alpha y \mathbb{1}_{[0,1]}(|y|)) \nu_i(dy),$$

where $\int_{-\infty}^{\infty} e^{\alpha y} y \mathbb{1}_{[0,1]}(|y|) \nu_i(dy) < \infty$. Note that the process $X^i(t)$ has the Lévy characteristics $(\alpha_i t, \sigma_i^2 t, \nu_i(dx))$. By $X^{(2)}(t)$ we denote the process which behaves in law like $X^i(t)$, when $J(t) = i$. Note that $(X^{(1)}(t), J(t))$ is a PDMP with extended generator

$$(\mathbf{A}^{(1)} f)(x, i) = \sum_{k=1}^N q_{ik} \int_{-\infty}^{\infty} (f(x+y, k) - f(x, i)) dG_i(y) = \sum_{k=1}^N q_{ik} \int_{-\infty}^{\infty} f(x+y, k) dG_i(y), \quad (5.28)$$

and domain $\mathcal{D}(\mathbf{A}^{(1)})$ consisting of absolutely continuous functions for which the above integrals are finite. By Çinlar *et al.* (1980, Th. 7.14, p. 211) and Jacod and Shiryaev (1987, Th. 2.42, p. 86), the process $X^i(t)$ has the following extended generator

$$\mathbf{A}^i f(x) = a_i f'(x) + \frac{\sigma_i}{2} f''(x) + \int_{-\infty}^{\infty} (f(x+y) - f(x) - \mathbb{1}_{[0,1]}(|y|) y f'(x)) \nu_i(dy),$$

with domain $\mathcal{C}^2(\mathbb{R}) \subset \mathcal{D}(\mathbf{A}^i)$. By Palmowski (1999, Th. 1), the process $(X^{(2)}(t), X^{(1)}(t), J(t))$ has the extended generator \mathbf{A} such that for $h(x, y, i) = g(x)p(y, i)$,

$$\mathbf{A}(gp)(x, y, i) = g(x) \mathbf{A}^{(1)} p(y, i) + p(y, i) \mathbf{A}^i g(x),$$

for $g \in \mathcal{C}^2(\mathbb{R}) \subset \mathcal{D}(\mathbf{A}^i)$ and $p \in \mathcal{D}(\mathbf{A}^{(1)})$ (that is, $\mathbf{A} = \mathbf{A}^{(1)} \oplus \mathbf{A}^i$ and $\mathcal{D}(\mathbf{A}^{(1)}) \otimes \mathcal{C}^2(\mathbb{R}) \subset \mathcal{D}(\mathbf{A})$).

Letting $\mathbf{Q} \circ \hat{\mathbf{G}}(\alpha) = (q_{ij}\hat{G}_i(\alpha))$, where $\hat{G}_i(\alpha) = \mathbb{E} \exp(\alpha U_i)$ is assumed to be finite, we define

$$\mathbf{F}(\alpha) = \mathbf{Q} \circ \hat{\mathbf{G}}(\alpha) + \text{diag}(\psi_1(\alpha), \dots, \psi_N(\alpha)) .$$

The Perron-Frobenius eigenvalue and the corresponding right eigenvector of $\mathbf{F}(\alpha)$ are denoted by $\lambda(\alpha)$ and $\mathbf{h}(\alpha)$, respectively. Note that $\lambda(\alpha)$ is real and $\mathbf{h}(\alpha)$ is positive. As a good function we propose $h(x, y, i) = g(x)f(y, i)$, where $g(x) = e^{\alpha x}$ and $p(y, i) = e^{\alpha y}h_i(\alpha)$. Then

$$(\mathbf{A}h)(x, y, i) = e^{\alpha(x+y)}(\psi_i(\alpha)h_i(\alpha) + (\mathbf{Q} \circ \hat{\mathbf{G}}(\alpha)\mathbf{h}(\alpha))_i) = \lambda(\alpha)h(x, y, i)e^{\alpha(x+y)} . \quad (5.29)$$

Proposition 5.6 *The process*

$$E^h(t) = e^{\alpha X(t) - \lambda(\alpha)t} \frac{h_{J(t)}(\alpha)}{h_{J(0)}(\alpha)}, \quad t \geq 0 , \quad (5.30)$$

is a mean-one martingale. Furthermore, the process $X(t)$, after the exponential change of measure, has the following characteristics:

$$\begin{cases} \tilde{q}_{ij} &= \frac{h_i(\alpha)}{h_i(\alpha)} q_{ij} \hat{G}_i(\alpha), \\ \tilde{G}_i(dx) &= \frac{e^{\alpha x}}{\hat{G}_i(\alpha)} G_i(dx), \\ \tilde{\sigma}_i &= \sigma_i, \\ \tilde{a}_i &= a + \alpha \sigma_i^2 - \int_{-\infty}^{\infty} |y| \mathbb{I}_{[0,1]}(|y|) (1 - e^{\alpha y}) \nu_i(dy), \\ \tilde{\nu}_i(dx) &= e^{\alpha x} \nu_i(dx). \end{cases} \quad (5.31)$$

Proof. By Lemma 3.1, $E^h(t)$ is a local martingale. Following Asmussen and Kella (2000) we have $\mathbb{E}E^h(t) = 1$ for all $t \geq 0$ and hence $E^h(t)$ is a true martingale. To prove (5.31) we write

$$\begin{aligned} \mathbf{A}(fh)(x, y, i) &= e^{\alpha(x+y)} h_i(\alpha) \sum_{k=1}^N \tilde{q}_{ik} \int_{-\infty}^{\infty} [f(x, y+z, k) - f(x, y, i)] d\tilde{G}_i(z) \\ &\quad + e^{\alpha(x+y)} h_i(\alpha) \tilde{\mathbf{A}}^i f(x, y, i) + e^{\alpha(x+y)} \sum_{k=1}^N q_{ik} f(x, y, i) h_k(\alpha) \hat{G}_i(\alpha) + e^{\alpha(x+y)} h_i(\alpha) f(x, y, i) \psi_i(\alpha) , \end{aligned}$$

which completes the proof in view of (5.29) and the formula: $\tilde{\mathbf{A}}f = h^{-1}(\mathbf{A}(fh) - f\mathbf{A}h)$.

□

For a one-dimensional Lévy process ($U_n^i = 0$ and $N = 1$), taking $h(x, y, i) = h(y) = e^{\alpha y}$, $h_i(\alpha) = 1$ and $\lambda(\alpha) = \psi(\alpha)$, we obtain the well known Wald martingale

$$\mathbb{E}^h(t) = e^{\alpha X(t) - \psi(\alpha)t} . \quad (5.32)$$

After the exponential change of measure, the Lévy process changes its Lévy characteristics from (a, σ, ν) to $(\tilde{a}, \sigma, \tilde{\nu})$, where

$$\begin{cases} \tilde{a} &= a + \alpha\sigma^2 - \int_{-\infty}^{\infty} |y| \mathbb{I}_{[0,1]}(|y|)(1 - e^{\alpha y})\nu(dy) \\ \tilde{\nu}(dx) &= e^{\alpha x} \nu_i(dx) ; \end{cases} \quad (5.33)$$

see K  chler and S  rensen (1997, Prop. 2.1.3, p. 11).

For a multi-dimensional Lévy process $(X_1(t), \dots, X_d(t))$ we take $h(x) = e^{\alpha^T x}$, where $\alpha, x \in \mathbb{R}^d$. This yields the martingale $E^h(t) = \prod_{k=1}^d e^{\alpha_k X_k(t) - \psi_k(\alpha_k)t}$; for further generalizations of this result see K  chler and S  rensen (1997, Th. A.10.1, p. 294), Jacod and M  min (1976, Th. 3.3, p. 13), and Jacod and Shiryaev (1987, Th. 3.24, p. 159).

Acknowledgment

The final draft of the paper was written while the second author was staying at the Department of Mathematical and Computing Sciences of Tokyo Institute of Technology. TR wishes to thank the Department and prof. Y. Takahashi for hospitality. ZP and TR are grateful to H. Schmidli, H.C. Gromoll and associate editor for their comments.

References

- [1] Asmussen, S. (1994) Busy period analysis, rare events and transient behaviour in fluid models. *Journal of Applied Math. and Stochastic Analysis* **7**(3), 269–299.
- [2] Asmussen, S. (1995) Stationary distribution for fluid models with or without Brownian noise. *Stochastic Models* **11**, 21–49.

- [3] Asmussen, S. (1998) *Stochastic simulation with a few towards stochastic processes*. Lecture Notes MaphySto no 2, Aarhus.
- [4] Asmussen, S. (2000) *Ruin Probabilities*. World Scientific Pub. Co., Singapore.
- [5] Asmussen, S. and Kella, O. (2000) Multi-dimensional martingale for Markov additive processes and its applications. *Adv. Appl. Probab.* **32(2)**, 376–380.
- [6] Çinlar, E., Jacod, J., Protter, P. and Sharpe, M.J. (1980) Semimartingales and Markov process. *Z. Wahrscheinlichkeitstheorieverw. Gebiete* **54**, 161–219.
- [7] Dassios, A. and Embrechts, P. (1989) Martingales and insurance risk. *Stochastic Models* **5(2)**, 181–217.
- [8] Davis, M.H.A. (1993) *Markov Models and Optimization*. Chapman and Hall, London.
- [9] Dellacherie C. and Meyer, P. (1982) *Probabilities and Potential B*. North-Holland Publishing Company, New York.
- [10] Doob, J.L. (1983) *Classical Potential Theory and Its Probabilistic Counterpart*, Springer, New York.
- [11] Dynkin, E.B. (1965) *Markov Processes*. **vol I**, 1965.
- [12] Ethier, H.J. and Kurtz, T.G. (1986) *Markov processes. Characterization and Convergence*. John and Wiley and Sons, Inc., New York.
- [13] Ethier, H.J. and Kurtz, T.G. (1993) Fleming-Viot processes in population genetic. *SIAM J. Control and Optimization* **31**, 345–386.
- [14] Gautam, N., Kulkarni, V., Palmowski, Z. and Rolski, T. (1999) Bounds for fluid models driven by semi-Markov inputs. *Prob.Eng.Inf.Sci.* **vol 18(4)**, 429–475.

- [15] Fukushima, M. and Stroock, D. (1986) Reversibility of solutions to martingale problems. *Probability, Statistical Mechanics, and Number Theory. Advanced in Mathematics Supplemental Studies*. vol. **9**, 107–123.
- [16] Itô, K. and Watanabe, S. (1965) Transformation of Markov processes by multiplicative functionals. *Ann. Inst. Fourier Grenoble* **15**, 13–30.
- [17] Jacod, J. and Mémin, J. (1976) Caractéristiques locales et conditions de continuité absolue pour les semi-martingales. *Z. Wahrscheinlichkeitstheorieverw. Gebiete* **35**, 1–37.
- [18] Jacod, J. and Shiryaev, A.N. (1987) *Limit Theorems for Stochastic Processes*. Springer-Verlag, Berlin, Heidelberg.
- [19] Karatzas, I. and Shreve, S.E. (1988) *Brownian Motion and Stochastic Calculus*. Springer-Verlag, New York.
- [20] Küchler, U. and Sørensen, M. (1997) *Exponential Families of Stochastic processes*. Springer-Verlag, New York.
- [21] Kulkarni, V. and Rolski, T. (1994) Fluid model driven by an Ornstein-Uhlenbeck process. *Prob.Eng.Inf.Sci.* **8**, 403–417.
- [22] Kunita, H. (1969) Absolute continuity of Markov processes and generators. *Nagoya Math. J.* **36**, 1–26.
- [23] Kunita, H. and Watanabe, T. (1963) Notes on transformations of Markov processes connected with multiplicative functionals. *Memoirs of the Faculty of Science, Kyushu University* **17**, ser. **A**, 181–191.
- [24] Liptser, R.S. and Shiryaev, A.N. (1986) *Teoriya martingalov*. "Nauka", Moskov.
- [25] Palmowski Z. (1999) Björk-Grandell model in a diffusion environment. Report no 119, Mathematical Institute, University of Wrocław, submitted for publication.

- [26] Palmowski, Z. and Rolski T. (1996) A note on martingales inequalities for fluid models. *Stochastic. Probab. Lett.* **31**, 13–21.
- [27] Palmowski, Z. and Rolski, T. (1998) Superposition of alternating on-off flows and a fluid model. *Ann.Appl.Prob.* **8(2)**, 524–541.
- [28] Parthasarathy. K.R. (1967) *Probability measures on metric spaces*. Academic Press, New York.
- [29] Protter, P. (1990) *Stochastic integration and differential equations. A new Approach*. Springer-Verlag, New York.
- [30] Revuz, D. and Yor, M. (1991) *Continuous martingales and Brownian Motion*, Springer–Verlag, Berlin.
- [31] Ridder, A. (1996) Fast simulation of Markov fluid models. *J. Appl. Prob.* **33(3)**, 786–804.
- [32] Ridder, A. and Walrand, J. (1992) Some large deviations results in Markov fluid models. *Prob. Eng. Inf. Sci.* **6**, 543–560.
- [33] Rogers, L.C.G. and Williams, D. (1987) *Diffusions, Markov Processes, and Martingales. Volume 2: Ito Calculus*. John Wiley and Sons, Inc., New York.
- [34] Rolski, T., Schmidli, H., Schmidt, V. and Teugles, J.L. (1999) *Stochastic processes for insurance and finance*. John Wiley and Sons, Inc., New York.
- [35] Schmidli, H. (1995) Cramér-Lundberg approximations for ruin functions of risk processes perturbed by diffusion. *Insurance: Mathematics and Economics* **16**, 135–149.
- [36] Schmidli, H. (1996) Lundberg inequalities for a Cox model with a piecewise constant intensity. *J. App.Prob.* **33**, 196–210.
- [37] Schmidli, H. (1997a). An extension to the renewal theorem and an application to risk theory. *Ann. Appl. Probab.* **7**, 121–133.

- [38] Schmidli, H. (1997b). Estimation of the Lundberg coefficient for a Markov modulated risk model. *Scandinavian Actuarial Journal*, 48–57.
- [39] Schwartz, A. and Weiss, A. (1995) *Large Deviations For Performance Analysis*. Chapman and Hall, London.
- [40] Stroock, D.W. (1987) *Lectures on Stochastic Analysis: Diffusion Theory*. Cambridge University Press.
- [41] Stroock, D.W. and Varadhan, S.R.S (1979) *Multidimensional diffusion processes*. Springer, New York.