

EDGEWORTH EXPANSIONS AND BOOTSTRAP FOR DEGENERATE VON MISES STATISTICS

BY

F. GÖTZE* (BIELEFELD) AND R. ZITIKIS** (VILNIUS)

Abstract. We prove Edgeworth expansions for degenerate von Mises statistics like the Beran, Watson, and Cramér-von Mises goodness-of-fit statistics. Furthermore, we show that the bootstrap approximation works up to an error of order $O(N^{-1/2})$ and that bootstrap based confidence regions attain a prescribed confidence level up to the order $O(N^{-1})$.

1. Introduction, main results, and some examples. Let $\{\Omega, \mathcal{A}, \mathbf{P}\}$ be a probability space, and let $X: \{\Omega, \mathcal{A}\} \rightarrow \{A, \mathcal{F}\}$ be a random variable. The distribution function of X will be denoted by \mathbf{F} . Furthermore, we shall denote by X_1, X_2, \dots independent copies of X .

In this paper we consider various approximations for the (degenerate) von Mises statistic

$$V_N \triangleq N^{-1} \sum_{j=1}^N \sum_{k=1}^N H(X_j, X_k),$$

where $H: A \times A \rightarrow \mathbf{R}$ is a symmetric measurable function which is assumed to be degenerate with respect to \mathbf{P} , that is

$$\mathbf{E}(H(X_1, X_2) | X_2) = 0 \text{ P-a.s.}$$

Furthermore, we assume

$$\mathbf{E}|H(X, X)| + \mathbf{E}H^2(X_1, X_2) < +\infty.$$

We use the notation

$$P_N(x) \triangleq \mathbf{P}\{V_N \leq x\}.$$

Our goal is to investigate the following three closely related problems for the statistic V_N :

* Research supported by the SFB 343 at Bielefeld.

** Research supported by the SFB 343 and Graduiertenkolleg at the University of Bielefeld.

- 1) A bootstrap approximation;
- 2) The bootstrap based coverage probabilities;
- 3) Edgeworth expansions.

The bootstrap version of V_N we use in this paper is defined (see [17]) as follows:

$$V_n^* \triangleq n^{-1} \sum_{j=1}^n \sum_{k=1}^n H^*(X_j^*, X_k^*),$$

$$H^*(x, y) \triangleq H(x, y) - E^* H(x, X_2^*) - E^* H(X_1^*, y) - E^* H(X_1^*, X_2^*),$$

where E^* denotes the expectation with respect to the empirical distribution function F^* given the sample X_1, \dots, X_N .

We shall use the notation

$$P_n^*(x) \triangleq P^*\{V_n^* \leq x\}.$$

The limit distribution of V_N , say P_∞ was described by von Mises [43], and it is the distribution of the random variable

$$V_\infty \triangleq EH(X, X) + \sum_{k=1}^{\infty} \lambda_k (G_k^2 - 1),$$

where $\lambda_1, \lambda_2, \dots, |\lambda_1| \geq |\lambda_2| \geq \dots$ are eigenvalues of the self-adjoint Hilbert-Schmidt operator $S: L_2(A, F) \rightarrow L_2(A, F)$ defined by

$$Sf \triangleq EH(\cdot, X)f(X) = \int H(\cdot, y)f(y)F(dy).$$

Here G_1, G_2, \dots are independent copies of a Gaussian random variable G with mean 0 and variance 1. See [31], [47], [49], and [19] for more details on this representation.

In the same way we may describe the distribution function P_∞^* of the random variable V_∞^* (\triangleq the weak limit, as $n \rightarrow +\infty$, of the statistic V_n^* when the random variables X_1, \dots, X_N are fixed).

If it is not stated otherwise, we shall always assume that the following holds:

The operator S has an infinite number of non-zero eigenvalues λ_j , and the random variables $H(X, X)$ and $H(X_1, X_2)$ have moments of all orders.

In the sequel we use $\|\cdot\|_\infty$ to denote the sup-norm.

THEOREM 1.1. *We have*

$$(1.1) \quad \|P_N - P_N^*\|_\infty = O_P(N^{-1/2}).$$

For the general theory of the bootstrap we refer to the papers [20], [13], [21], [2], [38], and the monograph [33].

Previous results on the CLT for degenerate von Mises statistics (which will be discussed below) show that the rate of convergence in the CLT for these

statistics might be of order $O(N^{-1+\varepsilon})$ for any $\varepsilon > 0$, and even of order $O(N^{-1})$ provided some smoothness conditions on the kernel H and the distribution function F are imposed. These facts are the main ingredients in the proof of the following theorem on the bootstrap based confidence regions; see [32], [8], and [33] on this topic. In the theorem below we use $P_N^*{}^{-1}(\alpha)$ to denote the α -th quantile of the function P_N^* .

THEOREM 1.2. *For each $\alpha \in (0, 1)$ and $\varepsilon > 0$*

$$(1.2) \quad P\{V_N \geq P_N^*{}^{-1}(\alpha)\} = 1 - \alpha + O(N^{-1+\varepsilon}), \quad N \rightarrow +\infty.$$

At this point a natural question arises: Does Theorem 1.2 hold with $\varepsilon = 0$? An inspection of the proof of Theorem 1.2 shows that (1.2) holds with $\varepsilon = 0$ if the distribution functions P_N and P_N^* have Edgeworth expansions of order $O(N^{-1})$ and $O_P(N^{-1-\delta})$ for some $\delta > 0$, respectively. In order to prove such a result we need to require more than moment conditions and assumptions on the limit distribution. As an example reflecting this fact we formulate the following result:

There exist infinitely differentiable and rapidly decreasing functions a_1, \dots, a_k such that the asymptotic expansion

$$(1.3) \quad P_N(x) = P_\infty(x) + a_1(x)N^{-1} + \dots + a_k(x)N^{-k} + R(x)$$

holds with a remainder term R satisfying, for any $\varepsilon > 0$ and some constant $c \geq 0$,

$$(1.4) \quad \sup_{x \in \mathbb{R}} |x|^m |R(x)| \leq \frac{c}{N^{k+1}} + c \sup_{0 \leq \mu \leq m} \int_{N^{1-\varepsilon} \leq |t| \leq N^{k+1}} \left| \left(\frac{d}{dt} \right)^\mu \mathbf{E} \exp\{itV_N\} \right| dt$$

for all $N \in \mathbb{N}$.

This result appeared in slightly different forms in [24], [26], and [27]; see also [7] and the survey paper [6].

Thus our further task is to investigate which assumptions on H and the distribution of the X yield bounds like

$$\sup_{|t| \geq \varrho(N)} |(d/dt)^m \mathbf{E} \exp\{itV_N\}| = O(N^{-L}), \quad N \rightarrow +\infty,$$

with some function $\varrho: N \rightarrow [0, +\infty)$ such that $\varrho(N) \rightarrow +\infty$ for $N \rightarrow +\infty$, and for every $L \geq 0$.

We assume that $\Lambda = \mathbb{R}^d$, and that the random variable X has a non-zero absolutely continuous component. This means that for some $\alpha \in (0, 1]$ the distribution function F of the X allows the representation

$$(1.5) \quad F = \alpha F_0 + (1 - \alpha)F_1,$$

where F_0 and F_1 are distribution functions, and F_0 is absolutely continuous. We use Y, Y_1, \dots, Y_N to denote i.i.d. \mathbb{R}^d -valued random variables having the distribution function F_0 .

We use $\mathcal{D}f(x)$ to denote the gradient of a function f at a point $x \in \mathbb{R}^d$. Also, if ξ is a random variable, let $\xi^{\#}$ denote a symmetrization of ξ .

THEOREM 1.3. *Let $m \in \mathbb{N} \cup \{0\}$ and*

$$(1.6) \quad \mathbb{E}|H(X_1, X_1)|^m + \mathbb{E}|H(X_1, X_2)|^m < +\infty.$$

Let B and C be some cubes in the space \mathbb{R}^d such that

$$(1.7) \quad \mathbb{P}\{Y \in B\} > 0, \quad \mathbb{P}\{Y \in C\} > 0.$$

Let Z_1, Z_2, \dots be independent \mathbb{R}^d -valued random variables having the distribution function $F_0\{\cdot|B\}$. Put

$$S_N(x) \triangleq N^{-1/2} \{ \tilde{H}(Z_1, x) + \dots + \tilde{H}(Z_N, x) \}.$$

Furthermore, assume that for every $\varepsilon > 0$ there exist numbers $\lambda \in \mathbb{R}$, $\kappa > 0$, and $\nu > 0$ such that $\lambda, \kappa, \nu \leq \varepsilon$ and such that for every $D \geq 0$ the following three conditions hold:

$$(1.8) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \{ \sup_{x \in C} \|\mathcal{D}S_N(x)\| \geq N^\lambda \} < +\infty;$$

$$(1.9) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \{ \sup_{x \in C} \|\mathcal{D}S_N(x)\| \geq 1/N^\kappa \} < +\infty;$$

$$(1.10) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \{ \sup_j \sup_{x, y \in C_j} \|\mathcal{D}S_N(x) - \mathcal{D}S_N(y)\| \geq 1/(2dN^\nu) \} < +\infty,$$

where $C_j, j = 1, \dots, [N^{\nu/d}]^d$ are the subcubes of the cube C such that $\text{Vol}(C_j) = \text{Vol}(C)/[N^{\nu/d}]^d$ for all $j = 1, \dots, [N^{\nu/d}]^d$, and $\|\cdot\|$ denotes the Euclidean norm in the space \mathbb{R}^d . Then, for every $\varepsilon > 0$ and $L \geq 0$,

$$(1.11) \quad \sup_{|t| \geq N^\varepsilon} |(d/dt)^m \mathbb{E} \exp\{itV_N\}| = O(N^{-L}), \quad N \rightarrow +\infty.$$

Some simple corollaries to Theorem 1.3 are given in Section 5 (see Corollaries 5.1–5.3 below). Let us note that in some special cases the validity of the bound (1.11) was investigated by Sadikova [48], Yurinskiĭ [55], van Zwet [58], Csörgő and Stachó [18], Götze [26], Zitikis [56], [57], Helmers [35], Bentkus et al. [7], etc. See also the survey paper by Bentkus et al. [6].

We shall now discuss applications of Theorems 1.1–1.3 to some goodness-of-fit statistics. Let $A = (0, 1)$, and let $X = U$, where U denotes a $(0, 1)$ -uniform random variable.

Beran's statistic B_N^2 . Let $b_0, b_1, \dots, \sum b_l^2 < +\infty$, denote the Fourier coefficients with respect to the basis $\{e^{i2\pi lx}, l \in \mathbb{Z}\}$ of a probability density f on the circle S^1 of unit circumference. If the function H is given by

$$(1.12) \quad H_1(x, y) \triangleq 2 \sum_{l=1}^{\infty} b_l^2 \cos 2\pi l(x-y),$$

then V_N is Beran's statistic B_N^2 ; see [9], [10]. Further investigations of the statistic B_N^2 , and more general ones as well, are done by Mardia [41], Giné [23], Prentice [46], and Baringhaus [4].

STATEMENT 1.1. Assume that the number of non-zero coefficients b_0, b_1, \dots is infinite and $\sum_{l=0}^{\infty} b_l^2 l^2 < +\infty$. Then

(i) Theorems 1.1 and 1.2 hold for Beran's statistic B_N^2 .

(ii) For every fixed $m \in N \cup \{0\}$, $L \geq 0$ and $\varepsilon > 0$ the bound (1.11) holds for all $t \in \mathbb{R}$ such that $|t| \geq N^\varepsilon$.

(iii) There exist infinitely differentiable and rapidly decreasing functions a_1, a_2, \dots such that, for every fixed k and $m \in N \cup \{0\}$, the asymptotic expansion (1.3) holds with the remainder term R satisfying

$$(1.13) \quad \sup_{x \in \mathbb{R}} |x|^m |R(x)| = O(N^{-k-1}), \quad N \rightarrow +\infty.$$

In the range $N^{1/2+\varepsilon} \leq |t| \leq c_1 N$ for some c_1 and any $\varepsilon > 0$, the bound (1.11) is given in [40].

Thus, in view of Statement 1.1 (iii) it follows that for Beran's statistic B_N^2 the bound of Theorem 1.2 holds with $\varepsilon = 0$ as well.

Watson's statistic W_N^2 . Here the function H is given by

$$H_2(x, y) \triangleq \mathbf{E}I(x, U)I(y, U),$$

where $I(x, v) \triangleq \mathbf{I}\{x \leq v\} - v - \mathbf{E}(\mathbf{I}\{x \leq U\} - U)$. Thus V_N is Watson's statistic W_N^2 (see [54]).

STATEMENT 1.2. (i) Theorems 1.1 and 1.2 hold for Watson's statistic W_N^2 .

(ii) For every fixed $m \in N \cup \{0\}$, $L \geq 0$ and $\varepsilon > 0$ the bound (1.11) holds for all $t \in \mathbb{R}$ such that $|t| \geq N^\varepsilon$.

(iii) There exist infinitely differentiable and rapidly decreasing functions a_1, a_2, \dots such that, for every fixed k and $m \in N \cup \{0\}$, the asymptotic expansion (1.3) holds with a remainder term R satisfying (1.13).

In the case $m = 0$ the result 1.2 (iii) was proved in [24].

Thus, in view of 1.2 (iii), the bound given in Theorem 1.2 for Watson's statistic W_N^2 is valid with $\varepsilon = 0$ as well.

The goodness-of-fit statistic $\omega_N^2(q)$. Let $q: (0, 1) \rightarrow [0, +\infty)$ be a measurable function such that $\int x(1-x)q(x)dx < +\infty$. Here the function H is given by

$$H_3(x, y) \triangleq \mathbf{E}J(x, U)J(y, U)q(U),$$

where $J(x, v) \triangleq \mathbf{I}\{x \leq v\} - v$, and V_N is the ω^2 -statistic; see [42] for an exhaustive review on ω^2 -statistic. Let us recall that in the case $q(x) = 1$ for all $x \in (0, 1)$ this is Cramér-von Mises' statistic, and in the case $q(x) = 1/x(1-x)$ for all $x \in (0, 1)$ it is Anderson-Darling's statistic.

STATEMENT 1.3. (i) *Theorems 1.1 and 1.2 hold for the Cramér–von Mises and Anderson–Darling statistics.*

(ii) *Let $m \in \mathbb{N} \cup \{0\}$ and assume that*

$$(1.14) \quad \mu(m) \triangleq \int_0^1 \left\{ \int_0^u sq(s) ds \right\}^m du + \int_0^1 \left\{ \int_u^1 sq(s) ds \right\}^m du < +\infty.$$

Furthermore, assume that there exists a non-empty interval $(\gamma, \delta) \subset (0, 1)$ such that

$$(1.15) \quad q(x) > 0 \quad \text{for all } x \in (\gamma, \delta),$$

and, for some numbers $\tau > 0$ and $c \geq 0$,

$$(1.16) \quad |q(x) - q(y)| \leq c|x - y|^\tau \quad \text{for all } x, y \in (\gamma, \delta).$$

Then, for every fixed $\varepsilon > 0$ and $L \geq 0$ the bound (1.11) holds for all $t \in \mathbb{R}$ such that $|t| \geq N^\varepsilon$.

(iii) *Fix $k, m \in \mathbb{N} \cup \{0\}$ and assume that $\mu(M) < +\infty$ for $M = k+2$ and $M = m$. If the assumptions (1.15) and (1.16) are satisfied, then there exist infinitely differentiable and rapidly decreasing functions a_1, a_2, \dots such that the asymptotic expansion (1.3) holds with the remainder R satisfying (1.13).*

In the case $p = 2$, Statement 1.3 (ii) improves Theorem 2.7 by Bentkus et al. [7] where the same bound (1.11) was proved under the condition $\sup\{|q'(x)|: x \in (\gamma, \delta)\} < +\infty$ (instead of (1.16)) and in the region $|t| \geq N^{1/2+\varepsilon}$ only. Let us note that if assumption (1.15) is valid, then there is an infinite number of non-zero eigenvalues λ_j ; see [7] and [6] for more details on this subject. Statement 1.3 (iii) improves the corresponding results for ω^2 -statistics given by Bentkus et al. [7].

In view of Statement 1.3 we claim that for the Cramér–von Mises and Anderson–Darling statistics the bound given in Theorem 1.2 is valid with $\varepsilon = 0$ as well.

2. Proof of Theorem 1.1. Let $S^*: L_2(A, \mathbb{F}^*) \rightarrow L_2(A, \mathbb{F}^*)$ be the operator defined by the formula

$$S^*f \triangleq \mathbf{E}^* H^*(\cdot, X^*) f(X^*) = \int H^*(\cdot, y) f(y) \mathbf{F}^*(dy).$$

Denote the eigenvalues of S^* by $\lambda_1^*, \lambda_2^*, \dots$, $|\lambda_1^*| \geq |\lambda_2^*| \geq \dots$. Given the random variables X_1, \dots, X_N , N fixed, V_n^* as n tends to infinity converges in distribution to the random variable

$$V_\infty^* \triangleq \mathbf{E}^* H^*(X^*, X^*) + \sum_{k=1}^{\infty} \lambda_k^* (G_k^2 - 1).$$

LEMMA 1.1. *For every $K \in \mathbb{N}$ and every $A \geq 0$ there exists a constant c such that*

$$(2.1) \quad \mathbf{P}\{|\lambda_k^*| \leq |\lambda_k| / \sqrt{2K}\} \leq cN^{-A}$$

for all $N \in \mathbb{N}$.

Proof. The idea of the proof is based on the proof of Lemma 4.1 by Bickel et al. [14]. Let us first note that without loss of generality we may assume $\lambda_K \neq 0$ and $A \geq 1$. Furthermore, let e_1, \dots, e_K be the eigenfunctions of the operator S corresponding to the eigenvalues $\lambda_1, \dots, \lambda_K$, and let $S_2: L_2(A, F) \rightarrow L_2(A, F)$ be the (self-adjoint and positive-definite) Hilbert-Schmidt operator corresponding to the kernel

$$(x, y) \mapsto \mathbb{E}H(x, X)H(X, y).$$

Then $\lambda_1^2, \dots, \lambda_K^2$ are the eigenvalues of the operator S_2 corresponding to the eigenfunctions e_1, \dots, e_K . Define

$$\mathcal{A} \triangleq \bigcup_{p=1}^K \{\|e_p\|_{F^*}^2 \geq 1/2\},$$

where $\|\cdot\|_{F^*}$ denotes the norm in the space $L_2(A, F^*)$. Since $\mathbb{E}e_p^2(X) = 1$, the quantity $1 - \mathbb{P}(\mathcal{A})$ does not exceed $\sum_{p=1}^K \mathbb{P}\{\mathbb{E}e_p^2(X) - \|e_p\|_{F^*}^2 \geq 1/2\}$, and therefore $1 - \mathbb{P}(\mathcal{A}) \leq cN^{-4}$. Thus we get the bound

$$(2.2) \quad \mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\} \leq \mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\} \cap \mathcal{A} + cN^{-4}.$$

Let δ_{pq} denote the Kronecker delta, and let

$$\mathcal{B} \triangleq \bigcap_{p=1}^K \bigcap_{q=1}^K \{ |N^{-1} \sum_{j=1}^N e_p(X_j)e_q(X_j) - \delta_{pq}| \leq 1/(2K) \}.$$

We clearly have

$$1 - \mathbb{P}(\mathcal{B}) \leq \sum_{p=1}^K \sum_{q=1}^K \mathbb{P}\{ |N^{-1} \sum_{j=1}^N e_p(X_j)e_q(X_j) - \delta_{pq}| \geq 1/2 \},$$

and, therefore, $1 - \mathbb{P}(\mathcal{B}) \leq cN^{-4}$. This bound together with (2.2) implies

$$(2.3) \quad \mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\} \leq \mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\} \cap \mathcal{A} \cap \mathcal{B} + cN^{-4},$$

and therefore our further task is to show that the bound (2.1) holds for the quantity $\mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\} \cap \mathcal{A} \cap \mathcal{B}$ instead of $\mathbb{P}\{|\lambda_K^*| \leq |\lambda_K|/\sqrt{2K}\}$.

Let us examine the event \mathcal{B} more closely. Assume that $c_1 e_1 + \dots + c_K e_K = 0$ in the space $L_2(A, F^*)$. This means that $\sum_{p=1}^K c_p e_p(X_j) = 0$ for all $j = 1, \dots, N$. Thus

$$\begin{aligned} 0 &= N^{-1} \sum_{j=1}^N \left| \sum_{p=1}^K c_p e_p(X_j) \right|^2 = \sum_{p=1}^K c_p^2 + \sum_{p=1}^K \sum_{q=1}^K c_p c_q \{ N^{-1} \sum_{j=1}^N e_p(X_j)e_q(X_j) - \delta_{pq} \} \\ &\geq \sum_{p=1}^K c_p^2 - (1/2K) \left\{ \sum_{p=1}^K |c_p|^2 \right\}^2 \geq \frac{1}{2} \sum_{p=1}^K c_p^2. \end{aligned}$$

Therefore $c_1 = \dots = c_K = 0$. Thus the functions e_1, \dots, e_K are linearly independent in the space $L_2(A, F^*)$.

Hence $e_p^\diamond \triangleq e_p / \|e_p\|_{F^*}$, $p = 1, \dots, K$, are well-defined (on the set \mathcal{A}) and linearly independent in the space $L_2(A, F^*)$ (on the set \mathcal{B}). Thus, $\mathcal{G} \triangleq \text{span}\{e_1^\diamond, \dots, e_K^\diamond\}$ is a K -dimensional subspace of the space $L_2(A, F^*)$.

Furthermore, let $S_2^* : L_2(A, F^*) \rightarrow L_2(A, F^*)$ be the (self-adjoint and positive-definite) Hilbert-Schmidt operator corresponding to the kernel

$$(x, y) \mapsto E^* H^*(x, X^*) H^*(X^*, y).$$

If e_1^*, \dots, e_K^* are the eigenfunctions of the operator S^* corresponding to the eigenvalues $\lambda_1^*, \dots, \lambda_K^*$, then $\lambda_1^{*2}, \dots, \lambda_K^{*2}$ are the eigenvalues of the operator S_2^* corresponding to the eigenfunctions e_1^*, \dots, e_K^* . Since \mathcal{G} is a K -dimensional subspace of the space $L_2(A, F^*)$, the following estimates hold:

$$(2.4) \quad \lambda_K^{*2} = \sup_{\mathcal{H}} \inf_{f \in \mathcal{H}} \frac{\|S_2^* f\|_{F^*}}{\|f\|_{F^*}} \geq \inf_{f \in \mathcal{G}} \frac{\|S_2^* f\|_{F^*}}{\|f\|_{F^*}} \geq \inf_{f \in \mathcal{G}} \frac{\langle S_2^* f, f \rangle_{F^*}}{\|f\|_{F^*}^2},$$

where \mathcal{H} denotes an arbitrary K -dimensional subspace of $L_2(A, F^*)$. Since $\delta_{pq} \lambda_p^2 = \langle S_2 e_p, e_q \rangle_F$, we have

$$(2.5) \quad \langle S_2^* e_p^\diamond, e_q^\diamond \rangle_{F^*} = \delta_{pq} \lambda_p^2 + \zeta_{pq},$$

where $\langle \cdot, \cdot \rangle_F$ and $\langle \cdot, \cdot \rangle_{F^*}$ denote the inner products in the spaces $L_2(A, F)$ and $L_2(A, F^*)$, respectively, and

$$\zeta_{pq} \triangleq \langle S_2 e_p, e_q \rangle_F - \langle S_2^* e_p^\diamond, e_q^\diamond \rangle_{F^*}.$$

Furthermore, since $f = \sum_{p=1}^K c_p e_p^\diamond$ for some $c \triangleq (c_1, \dots, c_K)$ and $\|e_p^\diamond\|_{F^*}^2 = 1$, we obtain $\|f\|_{F^*}^2 \leq K \|c\|^2$, where $\|c\|^2 \triangleq c_1^2 + \dots + c_K^2$ and, therefore,

$$\begin{aligned} \langle S_2^* f, f \rangle_{F^*} &= \langle (\delta_{pq} \lambda_p^2)_{p,q=1,\dots,K} c, c \rangle + \langle (\zeta_{pq})_{p,q=1,\dots,K} c, c \rangle \\ &\geq \lambda_K^2 \|c\|^2 - \max_{p,q=1,\dots,K} |\zeta_{pq}| \|c\|^2 \geq \{ \lambda_K^2 - \max_{p,q=1,\dots,K} |\zeta_{pq}| \} \|f\|_{F^*}^2 / K, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in the space R^K . Applying the just obtained estimate in the right-hand side of (2.4), we get

$$\lambda_K^{*2} \geq \{ \lambda_K^2 - \max_{p,q=1,\dots,K} |\zeta_{pq}| \} / K,$$

which implies

$$(2.6) \quad \begin{aligned} P(\{ |\lambda_K^*| \leq |\lambda_K| / \sqrt{2K} \} \cap \mathcal{A} \cap \mathcal{B}) &\leq P\{ \lambda_K^2 / (2K) \geq \{ \lambda_K^2 - \max_{p,q=1,\dots,K} |\zeta_{pq}| \} / K \} \\ &\leq P\{ \lambda_K^2 / 2 \leq \max_{p,q=1,\dots,K} |\zeta_{pq}| \}. \end{aligned}$$

Let us show that the right-hand side of (2.6) does not exceed cN^{-A} . This could be proved as follows: Write ξ_{pq} as the sum $\Delta_1 + \Delta_2 + \Delta_3$, where

$$\begin{aligned} \Delta_1 &\triangleq \langle S_2 e_p, e_q \rangle_{\mathbb{F}} - \langle S_2 e_p, e_q \rangle_{\mathbb{F}^*}, & \Delta_2 &\triangleq \langle S_2 e_p^\diamond, e_q^\diamond \rangle_{\mathbb{F}^*} \{ \|e_p\|_{\mathbb{F}^*} \|e_q\|_{\mathbb{F}^*} - 1 \}, \\ \Delta_3 &\triangleq \langle S_2 e_p^\diamond, e_q^\diamond \rangle - \langle S_2^* e_p^\diamond, e_q^\diamond \rangle_{\mathbb{F}^*}. \end{aligned}$$

One could easily prove that for every $j = 1, 2, 3$, every $A \geq 0$ and every positive constant c_1 , the quantities $\mathbb{P}\{| \Delta_j | \geq c_1\}$ do not exceed cN^{-A} . Since $\lambda_k^2 > 0$, this completes the proof of the lemma. ■

LEMMA 1.2. For every $A > 0$ there exists a constant $c > 0$ such that

$$\mathbb{P}\{\sqrt{N} \|P_\infty - P_\infty^*\|_\infty \geq a\} \leq cN^{-A} + ca^{-A}$$

for all $N \in \mathbb{N}$ and $a \geq 0$.

Proof. Because of Lemma 1.1 we only need to show that

(2.7)

$$\mathbb{P}\{\sqrt{N} \|P_\infty - P_\infty^*\|_\infty \geq a, |\lambda_k^*| \geq |\lambda_k|/\sqrt{2k}, k = 1, \dots, K\} \leq cN^{-A} + ca^{-A}$$

for a number K depending on A . In the following we estimate $\|P_\infty - P_\infty^*\|_\infty$ using Fourier's inversion formula. Let us first state some auxiliary results.

Let $L_2(A, \mathbb{F})$ denote the complex Hilbert space $L_2(A, \mathbb{C}, \mathbb{F})$ of all measurable functions $f: A \rightarrow \mathbb{C}$ such that

$$\|f\|_{\mathbb{F}}^2 \triangleq \mathbb{E} f(X) \bar{f}(X) = \int_A f \bar{f} d\mathbb{F} < +\infty.$$

The inner product in this space will be denoted by $\langle \cdot, \cdot \rangle_{\mathbb{F}}$. Also, let I denote the identity operator, and $\mathcal{T}_{\mathbb{F}}$ denote the trace operator in the space $L_2(A, \mathbb{F})$. Then

$$\mathbb{E} \exp\{itV_\infty\} = \exp\{T_t\},$$

where

$$T_t \triangleq it\mathbb{E}H(X, X) - 2 \int_0^t \mathcal{T}_{\mathbb{F}}\{S(I - 2ivS)^{-1}S\} v dv.$$

A similar representation holds for the quantity $\mathbb{E}^* \exp(itV_\infty^*)$.

Let us prove that there exists a constant $c \geq 0$, which might depend on K and λ_K only, and such that, for all $w \in [0, 1]$,

$$(2.8) \quad \Psi \triangleq |\exp\{wT_t^* + (1-w)T_t\}| \leq c(1+|t|)^{-K/2}.$$

Indeed, the representation

$$\mathcal{T}_{\mathbb{F}}\{S(I - 2ivS)^{-1}S\} = \mathcal{T}_{\mathbb{F}}\{S(I + 4v^2S^2)^{-1}S\} + 2iv\mathcal{T}_{\mathbb{F}}\{S(I + 4v^2S^2)^{-1}S^2\}$$

shows that

$$(2.9) \quad \operatorname{Re} \mathcal{F}_F \{S(I - 2ivS)^{-1}S\} = \sum_{k=1}^{\infty} \frac{\lambda_k^2}{1 + 4v^2 \lambda_k^2}.$$

A similar representation holds for the quantity $\operatorname{Re} \mathcal{F}_{F^*} \{S^*(I - 2ivS^*)^{-1}S^*\}$ as well. Therefore,

$$(2.10) \quad \Psi \leq \exp \left\{ -2w \int_0^t \sum_{k=1}^K \frac{\lambda_k^{*2}}{1 + 4v^2 \lambda_k^{*2}} v dv - 2(1-w) \int_0^t \sum_{k=1}^K \frac{\lambda_k^2}{1 + 4v^2 \lambda_k^2} v dv \right\}.$$

Since $\lambda_k^{*2} \geq \lambda_k^2/(2k)$ for all $k = 1, \dots, K$, the bound (2.10) implies $\Psi \leq c(1 + |t|)^{-K/2}$, which completes the proof of (2.8).

Starting from (2.8) we may use Fourier's inversion formula to bound the quantity $\|P_\infty - P_\infty^*\|_\infty$. We get

$$(2.11) \quad \|P_\infty - P_\infty^*\|_\infty \leq \Phi_1 \triangleq \int_{\mathbf{R}} \frac{1}{|t|} |\mathbf{E} \exp\{itV_\infty\} - \mathbf{E}^* \exp\{itV_\infty^*\}| dt.$$

Furthermore, the bound (2.8) implies

$$(2.12) \quad \Phi_1 = \int_0^1 \int_{\mathbf{R}} |\exp\{wT_t^* + (1-w)T_t\}| |t^{-1}\{T_t^* - T_t\}| dw dt \leq c\Phi_2,$$

where

$$\Phi_2 \triangleq \int_{\mathbf{R}} (1 + |t|)^{-K/2} |t^{-1}\{T_t^* - T_t\}| dt.$$

Let us prove the estimate

$$(2.13) \quad \mathbf{P}\{\sqrt{N} \Phi_2 \geq a\} \leq cN^{-A} + ca^{-A},$$

which clearly completes the proof of the lemma.

Let us introduce some additional notation. Denote the resolvent operator of S by $R(\cdot, S)$, that is $R(z, S) \triangleq (zI - S)^{-1}$ for any complex number z . Unless otherwise stated, we shall always assume $z = 1/(2iv)$. With these notations we may rewrite T_t as follows:

$$T_t = it \mathbf{E} H(X, X) + \int_0^t \mathbf{E} \langle iR(z, S)H(X, \cdot), H(X, \cdot) \rangle_F dv.$$

Also, a similar representation holds for the quantity T_t^* as well. Therefore,

$$t^{-1}\{T_t^* - T_t\} = ih_1 + t^{-1} \int_0^t h_2(v) dv,$$

where

$$h_1 \triangleq E^* H^*(X^*, X^*) - E H(X, X),$$

$$h_2(v) \triangleq E^* \langle iR(z, S^*) H^*(X^*, \cdot), H^*(X^*, \cdot) \rangle_{F^*} - E \langle iR(z, S) H(X, \cdot), H(X, \cdot) \rangle_F.$$

Consequently,

$$\Phi_2 \leq c|h_1| + \Delta[h_2],$$

where

$$\Delta[h_2] \triangleq \int_{\mathbf{R}} \int_0^t (1+|t|)^{-K/2} |t|^{-1} |h_2(v)| dv dt.$$

It is easy to see that $P\{\sqrt{N}|h_1| \geq a\} \leq cN^{-A} + ca^{-A}$. Therefore, in order to prove (2.13) we only need to show

$$(2.14) \quad P\{\sqrt{N} \Delta[h_2] \geq a\} \leq cN^{-A} + ca^{-A}.$$

Here we use the estimate $\Delta[h_2] \leq \Delta[h_3] + \Delta[h_4] + \Delta[h_5]$, where $h_3(v)$, $h_4(v)$, and $h_5(v)$ are the following three quantities, respectively:

$$E^* \langle R(z, S^*) H^*(X^*, \cdot), H^*(X^*, \cdot) \rangle_{F^*} - E^* \langle R(z, S^*) H(X^*, \cdot), H(X^*, \cdot) \rangle_{F^*},$$

$$E^* \langle R(z, S^*) H(X^*, \cdot), H(X^*, \cdot) \rangle_{F^*} - E^* \langle R(z, S) H(X^*, \cdot), H(X^*, \cdot) \rangle_{F^*},$$

$$E^* \langle R(z, S) H(X^*, \cdot), H(X^*, \cdot) \rangle_{F^*} - E \langle R(z, S) H(X, \cdot), H(X, \cdot) \rangle_F.$$

Some straightforward calculations show that

$$P\{\sqrt{N} \Delta[h_j] \geq a\} \leq cN^{-A} + ca^{-A}$$

for $j = 3, 4, 5$, which proves (2.14) and the lemma. ■

We will now prove Theorem 1.1. By Theorem (2.3) in [24], we have

$$(2.15) \quad \|P_N - P_\infty\|_\infty \leq c\beta_3^3 \lambda_{31}^{-9} / \sqrt{N},$$

where $\beta_3 \triangleq E|H(X, X)|^3 + E|H(X_1, X_2)|^3$. Let us note that the cited result also yields the bound

$$(2.16) \quad \|P_N^* - P_\infty^*\|_\infty \leq c(\beta_3^*)^3 (\lambda_{31}^*)^{-9} / \sqrt{N},$$

where β_3^* is similar to β_3 with the distribution function F replaced by the empirical distribution function F^* . Thus, the bounds (2.15) and (2.16) and Lemmas 1.1 and 1.2 together complete the proof of the theorem. ■

3. Proof of Theorem 1.2. For the proof of Theorem 1.2 we need the following lemma (cf. Lemma 1.1 by Beran [10]) several times:

LEMMA 3.1. For each $\alpha \in (0, 1)$ we have $P'_\infty(P_\infty^{-1}(\alpha)) > 0$.

Proof. Without loss of generality we may assume that $\lambda_1 > 0$. It is clear that the lemma follows from the following result:

There exists a point $x_0 \in [-\infty, +\infty)$ such that $P'_\infty(x) = 0$ for all $x \leq x_0$ and $P'_\infty(x) > 0$ for all $x > x_0$.

Let p and q denote, respectively, the densities of the random variables $\lambda_1(G_1^2 - 1)$ and $\sum_{k=2}^\infty \lambda_k(G_k^2 - 1)$. The density p vanishes for all $x \leq -\lambda_1$ and is positive for all $x > -\lambda_1$. Since the function P_∞ is continuous, the density q is not degenerated. Our claim follows from the representation $P'_\infty(x) = \int p(x-z)q(z)dz$ such that $P'_\infty(x) = 0$ for $x \leq x_0$ and $P'_\infty(x) > 0$ for all $x_0 < x < z$ for some $z \in \mathbf{R}$. The assumption $P'_\infty(z) = 0$ now leads to a contradiction with $P'_\infty(x) > 0$ for $x_0 < x < z$. ■

Without loss of generality we assume $\varepsilon > 0$ in the proof of Theorem 1.2 to be small (say $\varepsilon < 10^{-10}$) and $N \geq N_0$, where N_0 is a large fixed constant (which might depend on $\varepsilon > 0$ and $\alpha \in (0, 1)$).

We have

$$(3.1) \quad \mathbf{P}\{V_N \geq P_N^{*-1}(\alpha)\} = \mathbf{P}\{P_N^*(V_N) \geq \alpha\} = \mathbf{P}\{(P_N^* - P_\infty^*)(V_N) + P_\infty^*(V_N) \geq \alpha\}.$$

Let $\beta_0 = N^{-1+\varepsilon}$. Using the bound (2.5) of [24] (in the case $s = 4$) and Lemma 1.1 we get, for every $A \geq 0$,

$$\mathbf{P}\{\|P_N^* - P_\infty^*\|_\infty \geq \beta_0\} = O(N^{-4}).$$

Therefore, using a Slutsky argument (see Lemma 3 on p. 16 in [44]) in the right-hand side of (3.1), we get the bound (1.2) provided that

$$(3.2) \quad \mathbf{P}\{V_N \geq P_\infty^{*-1}(\gamma)\} = 1 - \gamma + O(N^{-1+\varepsilon})$$

with $\gamma = \alpha \pm \beta_0$.

Furthermore, since $P_\infty(P_\infty^{-1}(\gamma)) = \gamma$, we obtain

$$\begin{aligned} & \mathbf{P}\{V_N \geq P_\infty^{*-1}(\gamma)\} \\ &= 1 - \gamma + \mathbf{P}\{V_N - P_\infty^{*-1}(\gamma) + P_\infty^{-1}(\gamma) \geq P_\infty^{-1}(\gamma)\} - \mathbf{P}\{V_\infty \geq P_\infty^{-1}(\gamma)\}. \end{aligned}$$

Thus in order to show (3.2) it is enough to prove

$$(3.3) \quad \Delta \triangleq \|\mathbf{P}\{V_N - P_\infty^{*-1}(\gamma) + P_\infty^{-1}(\gamma) \geq \cdot\} - \mathbf{P}\{V_\infty \geq \cdot\}\|_\infty = O(N^{-1+\varepsilon}).$$

Write $q(f) \triangleq q_\gamma(f) \triangleq f^{-1}(\gamma)$, and let $h \triangleq P_\infty^* - P_\infty$.

Since $P'_\infty(P_\infty^{-1}(\gamma)) \geq \text{const}(P_\infty, \alpha) > 0$ (recall that $N \geq c_0$, where c_0 is a large fixed constant, and note that $P'_\infty(P_\infty^{-1}(\alpha)) > 0$), we infer that the quantity

$$q'(P_\infty)(h) \triangleq -h(P_\infty^{-1}(\gamma))/P'_\infty(P_\infty^{-1}(\gamma))$$

is well defined. We shall show at the end of the proof that on a set of probability close to 1 the quantity $q'(P_\infty)(h)$ is actually the directional derivative of the function q at the point P_∞ in the direction h .)

An application of Slutsky arguments shows that the quantity Δ does not exceed $\Delta_1 + \Delta_2 + \Delta_3$, where

$$\begin{aligned} \Delta_1 &\triangleq \|\mathbf{P}\{V_N - q'(P_\infty)(h) \geq \cdot\} - \mathbf{P}\{V_\infty \geq \cdot\}\|_\infty, \\ \Delta_2 &\triangleq \mathbf{P}\{|q(P_\infty^*) - q(P_\infty) - q'(P_\infty)(h)| \geq \beta_0\}, \\ \Delta_3 &\triangleq \sup_{x \in \mathbf{R}} \mathbf{P}\{x - \beta_0 \leq V_\infty \leq x + \beta_0\}. \end{aligned}$$

Using the fact that the Fourier transformation of P_∞ decreases rapidly, we easily arrive at the bound $\Delta_3 \leq c\beta_0 = O(N^{-1+\varepsilon})$. To prove the same bound for the quantity Δ_1 note first (see some details below) that, for any fixed real x ,

$$(3.4) \quad h(x) = N^{-1} \sum_{j=1}^N f(X_j, x) + r_N,$$

where f is a measurable function, the i.i.d. random variables $f(X_j, x)$ are centered and have finite moments of all orders. Furthermore, the remainder term r_N is such that for every $A \geq 0$ there exists a constant $c \geq 0$ such that

$$(3.5) \quad \mathbf{P}\{N|r_N| \geq a\} \leq cN^{-A} + ca^{-A}$$

for all $N \in \mathbf{N}$ and $a \geq 0$.

The asymptotic expansion (3.4) with the remainder term r_N as in (3.5) could be proved by using the Fourier transformation and (with slight modifications) following the lines of the proof of Lemma 1.2; we omit the details of the proof. Using Slutsky arguments together with the bounds (3.5) and $\Delta_3 \leq c\beta_0$ we obtain, for every $A \geq 0$,

$$(3.6) \quad \Delta_1 \leq \|\mathbf{P}\{V_N + N^{-1} \sum_{j=1}^N f(X_j, x_0)/P'_\infty(x_0) \geq \cdot\} - \mathbf{P}\{V_\infty \geq \cdot\}\|_\infty + O(N^{-1+\varepsilon}),$$

where $x_0 \triangleq P_\infty^{-1}(\gamma)$. The first summand on the right-hand side of (3.6) is of the order $O(N^{-1+\varepsilon})$; this fact follows from Corollary (3.20) by Götze [27] (see Example 3.7 therein as well). Thus we have proved $\Delta_1 = O(N^{-1+\varepsilon})$, and still have to show

$$(3.7) \quad \Delta_2 = O(N^{-1+\varepsilon}).$$

Let $\beta_1 \triangleq N^{-1/2+\varepsilon}$ and put $\mathcal{A} \triangleq \mathcal{A}_1 \cap \dots \cap \mathcal{A}_4$, where

$$\mathcal{A}_1 \triangleq \{|\lambda_K^*| \geq |\lambda_K|/\sqrt{2K}\}, \quad \mathcal{A}_{2+\kappa} \triangleq \{\|h^{(\kappa)}\|_\infty \leq \beta_1\}, \quad \kappa = 0, 1, 2$$

($h^{(\kappa)}$ denotes the κ -th derivative of the h ; $h^{(0)} \triangleq h$). We have already known that $1 - \mathbf{P}(\mathcal{A}_1) = O(N^{-A})$ (Lemma 1.1) and $1 - \mathbf{P}(\mathcal{A}_2) = O(N^{-A})$ (Lemma 1.2). Some slight modifications of the proof of Lemma 1.2 lead to the bound (valid for

every fixed $\kappa = 0, 1, 2, \dots$)

$$(3.8) \quad \mathbf{P}\{\sqrt{N} \|h^{(\kappa)}\|_{\infty} \geq a\} \leq cN^{-A} + ca^{-A},$$

which shows that $1 - \mathbf{P}(\mathcal{A}_{2+\kappa}) = O(N^{-A})$ for $\kappa = 1, 2$ (we omit the details since they are straightforward).

Therefore, we may restrict our analysis to the set \mathcal{A} only. On this set, for some constants c_1 and c_2 which do not depend upon N and $\tau \in [0, 1]$, the following two bounds hold:

$$H'_{\tau}(H_{\tau}^{-1}(\gamma)) \geq c_1 > 0 \quad \text{and} \quad |H''_{\tau}(H_{\tau}^{-1}(\gamma))| \leq c_2, \quad \text{where } H_{\tau} \triangleq P_{\infty} + \tau h.$$

This leads to the following Taylor expansion (with all the quantities well defined) for $q(P_{\infty}^*)$:

$$q(P_{\infty}^*) = q(P_{\infty}) + q'(P_{\infty})(h) + \int_0^1 (1-\tau) q''(H_{\tau})(h)^2 d\tau,$$

where

$$q''(H_{\tau})(h)^2 = 2h(H_{\tau}^{-1}(\gamma))h'(H_{\tau}^{-1}(\gamma))/H'_{\tau}(H_{\tau}^{-1}(\gamma))^2 \\ - h(H_{\tau}^{-1}(\gamma))^2 H''_{\tau}(H_{\tau}^{-1}(\gamma))/H'_{\tau}(H_{\tau}^{-1}(\gamma))^3.$$

After some tedious but elementary calculations, we obtain

$$\Delta_2 \leq \mathbf{P}\{\|h\|_{\infty} \|h'\|_{\infty} + \|h\|_{\infty}^2 \geq \beta_0\} + O(N^{-A}),$$

which together with the bound (3.8) completes the proof of (3.7), and of the theorem as well. ■

4. Proof of Theorem 1.3. In the proof we shall use the following lemma:

LEMMA 4.1. *Let $0 \leq a < b \leq 1$ be any numbers, and let g be a function differentiable on (a, b) such that $\text{sgn } g'(x) = \text{const}$ for all $x \in (a, b)$, and, for some numbers $\Phi > 0$ and Ψ ,*

$$\Phi \leq |g'(x)| \leq \Psi \quad \text{for all } x \in (a, b).$$

Furthermore, let h be a function integrable on (a, b) such that, for some $Y > 0$,

$$h(x) \geq Y \quad \text{for all } x \in (a, b).$$

Then, for all $\tau_0 \geq 0$ and all τ such that $|\tau| \geq \tau_0$,

$$(4.1) \quad I \triangleq \left| \int_a^b \exp\{itg(x)\} h(x) dx \right| \\ \leq \int_a^b h(x) dx - \frac{Y \min\{\Phi^3, 1\}}{48\Psi} \min\{\tau_0^2, 1\} (b-a)^3.$$

Proof. With some slight modifications we shall follow the lines of the proof of Lemma 2.1 by Götze and Hipp [30]. Without loss of generality we assume that $\text{sgn } g'(x) = 1$. Because of $h(g^{-1}(x)) \geq Y$ and $g'(g^{-1}(x)) \leq \Psi$ for all $x \in (g(a), g(b))$, we get

$$(4.2) \quad I \leq \left| \int_{g(a)}^{g(b)} \exp\{itx\} \left\{ \frac{h(g^{-1}(x))}{g'(g^{-1}(x))} - \frac{Y}{\Psi} \right\} dx \right| + \frac{Y}{\Psi} \left| \int_{g(a)}^{g(b)} \exp\{itx\} dx \right| \\ \leq \int_a^b h(x) dx - \frac{Y}{\Psi} \{g(b) - g(a)\} + \frac{Y}{\Psi} \left| \int_{g(a)}^{g(b)} \exp\{itx\} dx \right|.$$

Furthermore,

$$(4.3) \quad \left| \int_{g(a)}^{g(b)} \exp\{itx\} dx \right| = \{g(b) - g(a)\} \left| \frac{\sin v}{v} \right|,$$

where $v \triangleq \tau \{g(b) - g(a)\} / 2$. To estimate the quantity $|v^{-1} \sin v|$ on the right-hand side of (4.3) we use the bound $|x^{-1} \sin x| \leq 1 - \min\{d_0^2, 1\} / 12$ which is true for all $d_0 \geq 0$ and all $|x| \geq d_0$. Let us find a number d_0 such that $|v| \geq d_0$. It is clear that $|g(b) - g(a)| \geq \Phi(b - a)$. Therefore, $d_0 = \tau_0 \Phi(b - a) / 2$, and so we have

$$(4.4) \quad \left| \frac{\sin v}{v} \right| \leq 1 - \frac{\min\{\Phi^2, 1\}}{48} \min\{\tau_0^2, 1\} (b - a)^2.$$

Now taking the bounds (4.2)–(4.4) together, and using the bound $|g(b) - g(a)| \geq \Phi(b - a)$ once again, we get the lemma proved. ■

Let us note at the beginning of the proof of Theorem 1.3 that we do not specify the constants c in the text below; we only want to emphasize that all of them are non-random, do not depend on both N and t , and are non-negative. We subdivide the proof into several steps.

Step 1 (*reduction of the general case to the case $m = 0$*). Because of the moment condition (1.6), we interchange the signs of differentiation and integration in the quantity $(d/dt)^m \mathbf{E} \exp\{itV_N\}$. Then we write V_N^m as a multiple sum and use the fact that X_1, \dots, X_N are identically distributed. We get

$$|(d/dt)^m \mathbf{E} \exp\{itV_N\}| \leq cN^{2m} \mathbf{E} |\mathbf{E}^\ominus \exp\{itV_N\}|,$$

where \mathbf{E}^\ominus denotes the conditional expectation with respect to X_1, \dots, X_{N-2m} when all the other random vectors are fixed. Thus the theorem follows if

$$(4.5) \quad \mathbf{E} |\mathbf{E}^\ominus \exp\{itV_N\}| \leq cN^{-L}.$$

Step 2 (*using the absolutely continuous part*). Here we employ arguments used by Bikelis [16]. Without loss of generality we assume that $N \geq c_1$, where c_1 is a large constant (which might depend only on m and α). Write

$N_1 \triangleq N - 2m$ and fix X_{N-2m+1}, \dots, X_N . Because of decomposition (1.5), we have

$$(4.6) \quad \mathbf{E}^\ominus \exp\{itV_N\} = \int \exp\{itV_N\} (d\mathbf{F})^{N_1} \\ = \sum_{j=0}^{N_1} \binom{N_1}{j} \alpha^j (1-\alpha)^{N_1-j} \int \exp\{itV_N\} (d\mathbf{F}_0)^j (d\mathbf{F}_1)^{N_1-j}.$$

Now split the summation in (4.6) into two parts: $\sum' \triangleq$ the sum taken over all $j = 0, \dots, N_1$ such that $|j - \alpha N_1| < \sqrt{N_1} \log N_1$, and $\sum'' \triangleq$ the sum taken over all $j = 0, \dots, N_1$ such that $|j - \alpha N_1| \geq \sqrt{N_1} \log N_1$. A theorem by Bernstein [11] yields the bound

$$(4.7) \quad \sum_{\substack{j - \alpha N_1 \\ \geq 2x\sqrt{N_1\alpha(1-\alpha)}}} \binom{N_1}{j} \alpha^j (1-\alpha)^{N_1-j} \leq 2 \exp\{-x^2\}$$

for every $x \leq \sqrt{N_1\alpha(1-\alpha)}/4$. If we take $x = x_0$, where x_0 is the solution of the equation $2x\sqrt{\alpha(1-\alpha)} = \log N_1$, the bound (4.7) implies that the quantity

$$\sum'' \binom{N_1}{j} \alpha^j (1-\alpha)^{N_1-j}$$

does not exceed cN^{-L} . This bound and representation (4.6) together yield

$$(4.8) \quad |\mathbf{E}^\ominus \exp\{itV_N\}| \leq cN^{-L} + \sum' \binom{N_1}{j} \alpha^j (1-\alpha)^{N_1-j} \int |\exp\{itV_N\} (d\mathbf{F}_0)^j (d\mathbf{F}_1)^{N_1-j}|.$$

To estimate the second summand on the right-hand side of (4.8) we use the fact that $|j - \alpha N_1| < \sqrt{N_1} \log N_1$ (which implies that $j \geq [N_1\alpha/2]$). This fact and the estimate (4.8) show that the theorem will follow if we show

$$(4.9) \quad \mathbf{E} |\exp\{itV_N\} (d\mathbf{F}_0)^{[N_1\alpha/2]}| \leq cN^{-L}.$$

Step 3 (*reduction to the set $B \times C$*). The main idea of this step is based on arguments by Bikelis [16], Götze [24], and Bickel et al. [14]. Let $\beta \triangleq \mathbf{P}\{Y \in B\}$ and $\gamma \triangleq \mathbf{P}\{Y \in C\}$, and let

$$\mathbf{F}_2(\cdot) \triangleq \beta^{-1} \mathbf{P}\{Y \in \cdot \cap B\}, \quad \mathbf{F}_3(\cdot) \triangleq \gamma^{-1} \mathbf{P}\{Y \in \cdot \cap C\}.$$

Furthermore, let us put $N_2 \triangleq [N_1\alpha/4]$ and write

$$(4.10) \quad \int \exp\{itV_N\} (d\mathbf{F}_0)^{[N_1\alpha/2]} = \int \exp\{itV_N\} (d\mathbf{F}_0)^{N_2} (d\mathbf{F}_0)^{N_2} (d\mathbf{F}_0)^{[N_1\alpha/2] - 2N_2}.$$

In the first group of dF_0 on the right-hand side of (4.10) we decompose each F_0 as follows: $F_0 = \beta F_2 + (1 - \beta)$ ("some distribution function"), and in the second group of dF_0 on the right-hand side of (4.10) we decompose each F_0 as follows: $F_0 = \gamma F_3 + (1 - \gamma)$ ("some distribution function"). Now going along the lines of Step 2, we infer from (4.10) that in order to show (4.9) it is enough to prove the bound

$$I \triangleq \mathbf{E} \left| \int \exp \{itV_N\} (dF_2)^{[N_2\beta/2]} (dF_3)^{[N_2\gamma/2]} \right| \leq cN^{-L}.$$

Let Z, Z_1, Z_2, \dots , and W, W_1, W_2, \dots be independent random vectors having the distribution functions F_2 and F_3 , respectively. Using the symmetrization technique of the proof of Part 1 of Lemma (3.37) by Götze [24] we get

$$I \leq J \triangleq \mathbf{E} \left| \mathbf{E}^\oplus \exp \left\{ i \frac{2t}{N} \sum_{j=1}^{[N_2\beta/2]} \sum_{k=1}^{[N_2\gamma/2]} \{H(Z_j^*, W_k) - H(Z_j^{**}, W_k)\} \right\} \right|,$$

where Z_j^* and Z_j^{**} are independent and have the same distribution function F_2 . The \mathbf{E}^\oplus denotes the conditional expectation with respect to the random vectors $W_k, k = 1, \dots, [N_2\gamma/2]$. Therefore,

$$J = \mathbf{E} \left| \mathbf{E}^\oplus \exp \left\{ i \frac{2t}{N} \sum_{j=1}^{[N_2\beta/2]} \{H(Z_j^*, W) - H(Z_j^{**}, W)\} \right\} \right|^{[N_2\gamma/2]},$$

where \mathbf{E}^\oplus this time denotes the conditional expectation with respect to the random vector W . Thus in order to complete the proof of the theorem we need to show

$$(4.11) \quad J \leq cN^{-L}.$$

Step 4 (proof of the bound (4.11)). Let us rewrite the quantity J in a more convenient way for further calculations. Let us put $M \triangleq [N_2\beta/2]$, $\tau \triangleq 2t\sqrt{M}/N$, and let

$$\Delta_1 \triangleq \frac{1}{\gamma} \left| \int \exp \{i\tau S_M(x)\} p_0(x) dx \right|,$$

where p_0 denotes the density of F_0 . Then $J = \mathbf{E} \Delta_1^{[N_2\gamma/2]}$.

Fix an elementary event ω . Because of (1.8)–(1.10), we may assume without loss of generality that there exists a direction $k \in \{1, \dots, d\}$ and a point x such that

$$(4.12) \quad \sup_{x \in C} |\langle \mathcal{D}S_M(x), e_k \rangle| \leq M^\lambda,$$

$$(4.13) \quad |\langle \mathcal{D}S_M(x_0), e_k \rangle| \geq 1/(dM^k),$$

$$(4.14) \quad \sup_{x, y \in C_r} |\langle \mathcal{D}S_M(x) - \mathcal{D}S_M(y), e_k \rangle| \leq 1/(2dM^k),$$

where $C_r \in \{C_1, \dots, C_{[M^d]}\}$ is such that $x \in C_r$. Rewrite C_r as the Cartesian product $I_1 \times \dots \times I_d$ of intervals. Then, with dx_k^y denoting integration with respect to the variables different from x_k , we have

$$(4.15) \quad \begin{aligned} \Delta_1 &\leq \frac{1}{\gamma} \int_{C_r} p_0(x) dx + \frac{1}{\gamma} \left| \int_{C_r} \exp\{itS_M(x)\} p_0(x) dx \right| \\ &\leq \frac{1}{\gamma} \int_{C_r} p_0(x) dx + \frac{1}{\gamma} \int_{I_k} \left| \int \exp\{itS_M(x)\} p_0(x) dx_k \right| dx_k^y. \end{aligned}$$

To estimate the quantity $\left| \int_{I_k} \dots dx_k \right|$ we are going to use Lemma 4.1. For this, fix $x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_d$, and let $g(x_k) \triangleq S_M(x)$, $h(x_k) \triangleq p_0(x)$. Also, let a and b be numbers such that $I_k = (a, b)$. For every number $x \in C_r$, we have $g'(x_k) = \langle \mathcal{D}S_M(x), e_k \rangle$, which implies

$$(4.16) \quad |g'(x_k)| \geq |\langle \mathcal{D}S_M(x_0), e_k \rangle| - |\langle \mathcal{D}S_M(x) - \mathcal{D}S_M(x_0), e_k \rangle|.$$

Using (4.13) and (4.14) on the right-hand side of (4.16), we get $|g'(x_k)| \geq 1/(2dM^c)$. Therefore, we may choose

$$\Phi = 1/(2dM^c).$$

The same proof shows that for all $x \in C_r$, the sign of $g'(x_k)$ is the same (and equals $\text{sgn} \langle \mathcal{D}S_M(x_0), e_k \rangle$). For a number Ψ , as is easy to see from (4.12), we may use

$$\Psi = M^\lambda.$$

Since $\mathbf{P}\{Y \in C\} > 0$, there exists a cube $C_1 \subset C$ such that for some $c_1 > 0$ we have $p_0(x) > c_1$ for all $x \in C_1$. Thus (if necessary, replace the set C by C_1) for a number Y we may use

$$Y = c_1.$$

Thus, using Lemma 4.1, we get

$$\left| \int_{I_k} \exp\{itS_M(x)\} p_0(x) dx_k \right| \leq \int_{I_k} p_0(x) dx_k - cM^{-\lambda-3\kappa} \min\{\tau_0^2, 1\} (b-a)^3.$$

Hence, because of the bound (4.15), we have

$$(4.17) \quad \Delta_1 \leq 1 - c \text{Vol}(C_r) M^{-\lambda-3\kappa} \min\{\tau_0^2, 1\} (b-a)^2.$$

If we now use the bound $1 - x \leq \exp\{-x\}$ on the right-hand side of (4.17), then we obtain

$$J \leq \exp\{cM^{1-10\varepsilon} \min\{\tau_0^2, 1\}\}.$$

Take now $\tau_0^2 = M^{-1+10\varepsilon}(\log M)^2$ and note that the number $\varepsilon > 0$ may be taken as small as we want, say $\varepsilon \leq 10^{-10}$. Then the bound (4.11) follows immediately. This completes the proof. ■

5. Proof of the statements. Let us first prove Statement 1.3. To do that we give a special case of Theorem 1.3.

COROLLARY 5.1. *Let $m \in \mathbb{N} \cup \{0\}$ and*

$$(5.1) \quad \mathbb{E}|H(U_1, U_1)|^m + \mathbb{E}|H(U_1, U_2)|^m < +\infty.$$

Assume that there exists a non-empty rectangle $(\alpha, \beta) \times (\gamma, \delta) \subset (0, 1) \times (0, 1)$ such that for all $x \in (\alpha, \beta)$ the function $y \mapsto H(x, y)$ is differentiable on the interval (γ, δ) . Put $S_N \triangleq (T_1 + \dots + T_N)/\sqrt{N}$, where T_1, \dots, T_N are independent copies of the random function $T \triangleq \tilde{H}(\alpha + (\beta - \alpha)U, \cdot)$. Furthermore, assume that for every $\varepsilon > 0$ one may find numbers $\lambda \in \mathbb{R}, \kappa > 0$, and $\nu > 0$ such that $\lambda, \kappa, \nu \leq \varepsilon$ and such that for every $D \geq 0$ the following three conditions hold:

$$(5.2) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \left\{ \sup_{x \in (\gamma, \delta)} |S'_N(x)| \geq N^\lambda \right\} < +\infty;$$

$$(5.3) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \left\{ \|S'_N\|_{L_2(\gamma, \delta)} \leq 1/N^\kappa \right\} < +\infty;$$

$$(5.4) \quad \sup_{N \in \mathbb{N}} N^D \mathbb{P} \left\{ \sup_j \sup_{x, y \in I_j} |S'_N(x) - S'_N(y)| \geq 1/(2N^\nu) \right\} < +\infty,$$

where $I_j \subset (\gamma, \delta), j = 1, \dots, [N^\nu]$, are subintervals of the interval (γ, δ) such that $\text{Vol}(I_j) = (\delta - \gamma)/[N^\nu]$. Then, for every $\varepsilon > 0$ and $L \geq 0$,

$$\sup_{t: |t| \geq N^\varepsilon} |(d/dt)^m \mathbb{E} \exp\{itV_N\}| = O(N^{-L}), \quad N \rightarrow +\infty.$$

Proof. The corollary is an easy consequence of Theorem 1.3. Let us note only that the condition (1.9) follows from (5.3) by using the inequality

$$(5.5) \quad \|S'_N\|_{L_2(\gamma, \delta)} \leq \|S'_N\|_{L_\infty(\gamma, \delta)} \sqrt{\delta - \gamma}. \quad \blacksquare$$

Also, to prove Statement 1.3 we need the following lemma which is a simple consequence of Corollary 1.2 by Bentkus [5].

LEMMA 5.1. *Let \mathcal{G} be a centered Gaussian $L_2(\gamma, \delta)$ -valued random variable the covariance of which is the Hilbert-Schmidt operator $L_2(\gamma, \delta) \rightarrow L_2(\gamma, \delta)$ corresponding to the kernel*

$$(x, y) \mapsto \mathbb{E} \left\{ \frac{d}{dx} H(\alpha + (\beta - \gamma)U, x) \frac{d}{dy} H(\alpha + (\beta - \gamma)U, y) \right\} \\ - \mathbb{E} \left\{ \frac{d}{dx} H(\alpha + (\beta - \gamma)U, x) \right\} \mathbb{E} \left\{ \frac{d}{dy} H(\alpha + (\beta - \gamma)U, y) \right\}.$$

If, for some $l > 2$,

$$(5.6) \quad \mathbb{E} \left\{ \int_\gamma^\delta \left| \frac{d}{dx} H(\alpha + (\beta - \alpha)U, x) \right|^2 dx \right\}^{1/2} < +\infty,$$

and the random variable \mathcal{G} is not concentrated in any finite-dimensional subspace of $L_2(\gamma, \delta)$, then the condition (5.3) is satisfied.

Proof of Statement 1.3. (i) This is an immediate consequence of Theorem 1.1.

(ii) Let us verify the conditions of Corollary 5.1 with the function H_3 when $d = 1$, $\alpha = 0$, and $\beta = 1$. The equivalence of the moment conditions (5.1) and (1.14) is easy to prove. Thus let us verify (5.2)–(5.4). Note first that

$$(5.7) \quad S'_N(x) = \frac{1}{\sqrt{N}} \sum_{j=1}^N \frac{d}{dx} \tilde{H}_3(U_j, x) = -\tilde{\mathcal{E}}_N(x)q(x),$$

where $\tilde{\mathcal{E}}_N$ denotes the uniform empirical process. Let us look at the condition (5.2). It is clear that without loss of generality we may assume

$$(5.8) \quad \sup\{q(x) : x \in (\gamma, \delta)\} < +\infty.$$

Thus, because of (5.8) and Lemma 2.3 by Stute [53], we obtain

$$(5.9) \quad \mathbf{P}\left\{\sup_{x \in (\gamma, \delta)} |\tilde{\mathcal{E}}_N(x)q(x)| \geq N^\lambda\right\} \leq \mathbf{P}\left\{c \sup_{x \in (\gamma, \delta)} |\tilde{\mathcal{E}}_N(x)| \geq N^\lambda\right\} \leq cN^{-D},$$

which completes the verification of (5.2).

The condition (5.3) is satisfied because of Lemma 2.1 and (5.1); compare the discussion concerning the infinite-dimensionality of the weighted Brownian bridge given just after Theorem 1.4 in [6].

Let us now show that (1.16) implies (5.4). Write

$$\Delta \triangleq \mathbf{P}\left\{\sup_j \sup_{x, y \in I_j} |\tilde{\mathcal{E}}_N(x)q(x) - \tilde{\mathcal{E}}_N(y)q(y)| \geq 1/(2N^\kappa)\right\}.$$

The assumptions (1.16) and (5.8) imply that, for some constant $c_2 > 0$,

$$(5.10) \quad \Delta \leq \mathbf{P}\left\{\sup_j \sup_{x, y \in I_j} |\tilde{\mathcal{E}}_N(x) - \tilde{\mathcal{E}}_N(y)| + \sup_j \sup_{x, y \in I_j} |\tilde{\mathcal{E}}_N(x)| N^{-\nu\tau} \geq c_2 N^{-\kappa}\right\}.$$

Using the bound (5.9) with $q(x) = 1$ on the right-hand side of (5.10), for a constant $c > 0$ we get

$$(5.11) \quad \begin{aligned} \Delta &\leq \mathbf{P}\left\{\sup_j \sup_{x, y \in I_j} |\tilde{\mathcal{E}}_N(x) - \tilde{\mathcal{E}}_N(y)| + N^{\lambda - \nu\tau} \geq cN^{-\kappa}\right\} + cN^{-D} \\ &\leq \mathbf{P}\left\{\sup_j \sup_{x, y \in I_j} |\tilde{\mathcal{E}}_N(x) - \tilde{\mathcal{E}}_N(y)| \geq cN^{-\kappa}\right\} + cN^{-D} \end{aligned}$$

if $\kappa < \nu\tau - \lambda$. Using Lemma 2.4 by Stute [53] or Inequality 3.2 by Shorack and Wellner [51] it follows that for small numbers $\nu > 0$ and $\kappa > 0$ the right-hand side of (5.11) does not exceed cN^{-D} . Let us note also that the numbers $\lambda > 0$, $\nu > 0$ and $\kappa > 0$ could be chosen arbitrarily small (but satisfying the condition $\kappa < \nu\tau - \lambda$).

(iii) The result is a direct consequence of part (ii) of this statement; see Section 3 (or Section 2) in [7] for more details. ■

Let us now prove Statement 1.2. For this we formulate another special case of Theorem 1.3.

COROLLARY 5.2. Assume that

$$(5.12) \quad \sup_{x,y \in (0,1)} \left| \frac{d}{dy} H(x, y) \right| < +\infty.$$

Let us put $S'_N \triangleq (T_1 + \dots + T_N)/\sqrt{N}$, where T_1, \dots, T_N are independent copies of the random function $T \triangleq \tilde{H}(U, \cdot)$. Furthermore, assume that for every $\varepsilon > 0$ one may find numbers $\kappa > 0$ and $\nu > 0$ such that $\kappa, \nu \leq \varepsilon$ and such that for every $D \geq 0$ the following condition holds:

$$(5.13) \quad \sup_{N \in \mathbb{N}} N^D \mathbf{P} \left\{ \sup_j \sup_{x,y \in I_j} |S'_N(x) - S'_N(y)| \geq 1/(2N^\kappa) \right\} < +\infty,$$

where $I_j \subset (0, 1)$, $j = 1, \dots, [N^\nu]$, are subintervals of the interval $(0, 1)$ such that $\text{Vol}(I_j) = 1/[N^\nu]$. Furthermore, let \mathcal{G} be a centered Gaussian $L_2(0, 1)$ -valued random variable the covariance of which is the Hilbert-Schmidt operator $L_2(0, 1) \rightarrow L_2(0, 1)$ corresponding to the kernel

$$(5.14) \quad (x, y) \mapsto \mathbf{E} \left\{ \frac{d}{dx} H(U, x) \frac{d}{dy} H(U, y) \right\} - \mathbf{E} \left\{ \frac{d}{dx} H(U, x) \right\} \mathbf{E} \left\{ \frac{d}{dy} H(U, y) \right\}.$$

Assume that the random element \mathcal{G} is not concentrated in any finite-dimensional subspace of $L_2(0, 1)$. Then, for every $m \in \mathbb{N} \cup \{0\}$, $\varepsilon > 0$ and $L \geq 0$,

$$\sup_{t: |t| \geq N^\varepsilon} |(d/dt)^m \mathbf{E} \exp\{itV_N\}| = O(N^{-L}), \quad N \rightarrow +\infty.$$

Proof. We shall show that Corollary 5.1 implies the result. Take $\alpha = \gamma = 0$ and $\beta = \delta = 1$. Then, for every $j = 1, \dots, [N^\nu]$, choose a non-random point $y_j \in I_j$. Using Slutsky's arguments, we get

$$(5.15) \quad \begin{aligned} \Delta &\triangleq \mathbf{P} \left\{ \sup_{x \in (0,1)} |S'_N(x)| \geq N^\lambda \right\} = \mathbf{P} \left\{ \sup_j \sup_{x \in I_j} |S'_N(x)| \geq N^\lambda \right\} \\ &\leq \mathbf{P} \left\{ \sup_j \sup_{x,y \in I_j} |S'_N(x) - S'_N(y)| \geq 1/(2N^\kappa) \right\} + \mathbf{P} \left\{ \sup_j |S'_N(y_j)| + 1/(2N^\kappa) \geq N^\lambda \right\}. \end{aligned}$$

Because of (5.13), the first summand on the right-hand side of inequality (5.15) does not exceed cN^{-D} . Therefore, $\Delta \leq cN^{-D}$ follows if

$$(5.16) \quad \mathbf{P} \left\{ \sup_j |S'_N(y_j)| \geq N^\lambda/2 \right\} \leq cN^{-D}.$$

But the bound (5.16) (and (5.2) as well) are consequences of Markov's inequality and the bound $E|S'_N(y_k)|^p \leq c$ which holds, because of the assumption (5.12), for all $k = 1, \dots, [N^\nu]$ and all $D \geq 0$, where the constant $c \geq 0$ does not depend on N . Furthermore, because of Lemma 5.1 the assumption (5.3) is satisfied as well. This remark completes the proof of the theorem. ■

Proof of Statement 1.2. (i) This is an easy consequence of Theorem 1.1.

(ii) The proof is almost the same as that of Statement 1.3; use Corollary 5.2 instead of Corollary 5.1.

(iii) This is a consequence of part (ii); use results from Section 3 of [7]. ■

Finally, as a simple consequence of Theorem 1.3 we have

COROLLARY 5.3. Assume that

$$(5.17) \quad \sup_{x, y \in (0, 1)} \left| \left(\frac{d}{dy} \right)^2 H(x, y) \right| < +\infty.$$

Furthermore, let \mathcal{G} be a centered Gaussian $L_2(0, 1)$ -valued random variable with covariance being the Hilbert-Schmidt operator $L_2(0, 1) \rightarrow L_2(0, 1)$ corresponding to the kernel defined by (5.14). Assume that the random element \mathcal{G} is not concentrated in any finite-dimensional subspace of $L_2(0, 1)$. Then for every $m \in \mathbb{N} \cup \{0\}$, $\varepsilon > 0$ and $L \geq 0$,

$$\sup_{t: |t| \geq N^\varepsilon} |(d/dt)^m E \exp\{itV_N\}| = O(N^{-L}), \quad N \rightarrow +\infty.$$

Proof. This corollary follows from Corollary 5.2. Note that the assumption (5.12) holds because of (5.17). Furthermore, using the bound

$$\sup_{N \in \mathbb{N}} N^D \sup_{x, y \in I_j} |S'_N(x) - S'_N(y)| \leq \|S''_N(\cdot)\|_{L_2(I_j)} / \sqrt{[N^\nu]},$$

we see that the left-hand side of (5.13) does not exceed

$$(5.18) \quad \mathbb{P}\{\|S''_N(\cdot)\|_{L_2(I_j)} \geq \sqrt{[N^\nu]} / (2N^\kappa)\}.$$

If $\nu > 2\kappa$, then for every $D \geq 0$ the quantity (5.18) does not exceed cN^{-D} because of the Markov inequality and (5.17). This completes the proof of the theorem. ■

Proof of Statement 1.1. (i) This is an easy consequence of Theorem 1.1.

(ii) Because of $\sum_{l=0}^{\infty} b_l^2 l^2 < +\infty$ we have

$$(5.19) \quad \frac{d^2}{dx dy} H_1(x, y) = -2(2\pi)^2 \sum_{l=1}^{\infty} b_l^2 l^2 \cos 2\pi l(x-y)$$

for all $x, y \in (0, 1)$, which shows that the condition (5.14) holds. The assumption on infinite-dimensionality of the corresponding Gaussian random variable is

satisfied because there is an infinite number of non-zero coefficients b_0, b_1, \dots . Thus, Corollary 5.3 implies the desired result.

(iii) Because of (5.19) we have

$$B_N^2 \triangleq \frac{1}{N} \sum_{j=1}^N \sum_{k=1}^N H(U_j, U_k) = \int_0^1 \int_0^1 \mathcal{E}_N(x) \mathcal{E}_N(y) \frac{d^2}{dx dy} H(x, y) dx dy,$$

which shows that $B_N^2 = \pi_2(\mathcal{E}_N, \mathcal{E}_N)$, where π_2 is a polynomial of degree 2 in the Hilbert space $L_2(0, 1)$. Therefore, results by Bentkus et al. [7] might be used to get the theorem proved. (In the case $m = 0$ one may use general results for von Mises statistics by Götze [24], [26].) ■

Acknowledgement. The authors would like to thank V. Bentkus for interesting discussions and for several suggestions which led to simplifications of some of the proofs.

REFERENCES

- [1] T. W. Anderson and D. A. Darling, *Asymptotic theory of certain "goodness of fit" criteria based on stochastic processes*, Ann. Math. Statist. 23 (1952), pp. 193–212.
- [2] K. B. Athreya, *Strong law for the bootstrap*, Statist. Probab. Lett. 1 (1983), pp. 147–150.
- [3] – M. Ghosh, Y. L. Low and P. K. Sen, *Laws of large numbers for bootstrapped U-statistics*, J. Statist. Plann. Inference 9 (1984), pp. 185–194.
- [4] L. Baringhaus, *Testing for spherical symmetry of a multivariate distribution*, Ann. Statist. 19 (1991), pp. 899–917.
- [5] V. Bentkus, *On concentration functions of sums of independent random elements with values in a Banach space*, Litovsk. Mat. Sb. 14 (1986), pp. 32–39.
- [6] – F. Götze, V. Paulauskas and A. Račkauskas, *The accuracy of Gaussian approximation in Banach spaces*, SFB 343 "Diskrete Strukturen in der Mathematik", preprint 90–100, Universität Bielefeld, 1990.
- [7] V. Bentkus, F. Götze and R. Zitikis, *Asymptotic expansions in the integral and local limit theorems in Banach spaces with applications to ω -statistics*, J. Theoret. Probab. 6 (1993), pp. 727–780.
- [8] R. Beran, *Prepivoting to reduce level error of confidence sets*, Biometrika 74 (1987), pp. 457–468.
- [9] R. J. Beran, *Testing for uniformity on a compact homogeneous space*, J. Appl. Probab. 5 (1968), pp. 177–195.
- [10] – *Asymptotic theory of a class of tests for uniformity of a circular distribution*, Ann. Math. Statist. 40 (1969), pp. 1196–1206.
- [11] S. N. Bernstein, *On modification of Čebyšev inequality and an error in Laplace's formula*, Učen. Zap. Naučno-Issled. Kafedr. Ukrain. Otd. Mat. 1 (1924), pp. 38–49. Reprinted in: *Collected Works*, Vol. IV, Nauka, Moscow 1964, pp. 71–79 (in Russian).
- [12] R. N. Bhattacharya and Rao R. Ranga, *Normal Approximation and Asymptotic Expansions*, Wiley, New York 1976.
- [13] P. J. Bickel and D. Freedman, *Some asymptotic theory for the bootstrap*, Ann. Statist. 9 (1981), pp. 1196–1217.
- [14] P. J. Bickel, F. Götze and W. R. van Zwet, *The Edgeworth expansion for U-statistics of degree two*, ibidem 14 (1986), pp. 1463–1484.

- [15] A. Bikelis, *Remainder terms in asymptotic expansions for characteristic functions and their derivatives*, Selected Transl. Math. Statist. Probab. 11 (1973), pp. 149–162.
- [16] — *Asymptotic expansions for the densities and distributions of sums of independent identically distributed random vectors*, *ibidem* 13 (1973), pp. 213–234.
- [17] J. Bretagnolle, *Lois limites du bootstrap de certaines fonctionnelles*, Ann. Inst. H. Poincaré Sect. B 19 (1983), pp. 281–296.
- [18] S. Csörgő and L. Stachó, *A step toward asymptotic expansions for the Cramér–von Mises statistic*, in: *Analytic Function Methods in Probability Theory*, North-Holland, 1980, pp. 45–67.
- [19] E. B. Dynkin and A. Mandelbaum, *Symmetric statistics, Poisson point processes, and multiple Wiener integrals*, Ann. Statist. 31 (1983), pp. 739–745.
- [20] B. Efron, *Bootstrap methods: Another look at the jackknife*, *ibidem* 7 (1979), pp. 1–26.
- [21] — *The Jackknife, the Bootstrap and Resampling Plans*, SIAM, Philadelphia 1982.
- [22] A. A. Filippova, *Mises theorem of the asymptotic behavior of functionals of empirical distribution function and its statistical applications*, Theory Probab. Appl. 7 (1962), pp. 26–60.
- [23] E. M. Giné, *Invariant tests for uniformity on compact Riemannian manifolds based on Sobolev norms*, Ann. Statist. 3 (1975), pp. 1243–1266.
- [24] F. Götze, *Asymptotic expansions for bivariate von Mises functionals*, Z. Wahrscheinlichkeitstheorie verw. Gebiete 50 (1979), pp. 333–355.
- [25] — *On Edgeworth expansions in Banach spaces*, Ann. Probab. 9 (1981), pp. 852–859.
- [26] — *Expansions for von Mises functionals*, Z. Wahrscheinlichkeitstheorie verw. Gebiete 65 (1984), pp. 599–625.
- [27] — *Asymptotic expansion in functional limit theorem*, J. Multivariate Anal. 16 (1985), pp. 1–20.
- [28] — *On the rate of convergence in the central limit theorem in Banach spaces*, Ann. Probab. 14 (1986), pp. 922–942.
- [29] — *Edgeworth expansions in functional limit theorems*, *ibidem* 17 (1989), pp. 1602–1634.
- [30] — and C. Hipp, *On the validity of Edgeworth-expansions*, SFB 343 “Diskrete Strukturen in der Mathematik”, preprint 92-045, Universität Bielefeld, 1992.
- [31] G. G. Gregory, *Large sample theory for U-statistics and tests of fit*, Ann. Statist. 5 (1977), pp. 110–123.
- [32] P. Hall, *On the bootstrap and confidence intervals*, *ibidem* 14 (1986), pp. 1431–1452.
- [33] — *The Bootstrap and Edgeworth Expansions*, Springer, New York 1992.
- [34] R. Helmers, *Edgeworth expansions for linear combinations of order statistics*, Math. Center Tracts, Vol. 103, Amsterdam 1982.
- [35] — *A local limit theorem for L-statistics*, CWI Preprint (1991).
- [36] — *On the Edgeworth expansion and the bootstrap approximation for a studentized U-statistic*, Ann. Statist. 19 (1991), pp. 470–484.
- [37] — P. Janssen and R. Serfling, *Berry–Esseen and bootstrap results for generalized L-statistics*, Scand. J. Statist. 17 (1990), pp. 65–77.
- [38] D. V. Hinkley, *Bootstrap methods (With discussion)*, J. Roy. Statist. Soc. Ser. B 50 (1988), pp. 321–337.
- [39] T. Inglot, W. C. M. Kallenberg and T. Ledwina, *Asymptotic behaviour of some bilinear functionals of the empirical process*, Memorandum of Universiteit Twente 1031 (1992), pp. 1–27.
- [40] V. S. Koroliuk and Yu. V. Borovskich, *Asymptotic Analysis of Distributions of Statistics* (in Russian), Naukova Dumka, Kiev 1984.
- [41] K. V. Mardia, *Statistics of Directional Data*, Academic Press, New York 1972.
- [42] G. V. Martynov, *Omega-square Criterion* (in Russian), Nauka, Moscow 1979.

- [43] R. von Mises, *On the asymptotic distribution of differentiable statistical functions*, Ann. Math. Statist. 18 (1947), pp. 309–348.
- [44] V. V. Petrov, *Sums of Independent Random Variables*, Springer, Berlin 1975.
- [45] A. N. Pettitt and M. A. Stephens, *Modified Cramér–von Mises statistics for censored data*, Biometrika 63 (1976), pp. 291–298.
- [46] M. J. Prentice, *On invariant tests of uniformity for directions and orientations*, Ann. Statist. 6 (1978), pp. 169–176.
- [47] H. Rubin and R. A. Vitale, *Asymptotic distribution of symmetric statistics*, ibidem 8 (1980), pp. 165–170.
- [48] S. M. Sadikova, *Some inequalities for characteristic functions*, Probab. Theory Appl. 11 (1966), pp. 441–447.
- [49] R. J. Serfling, *Approximation Theorems of Mathematical Statistics*, Wiley, New York 1980.
- [50] G. R. Shorack and J. A. Wellner, *Limit theorems and inequalities for the uniform empirical process indexed by intervals*, Ann. Probab. 10 (1982), pp. 639–652.
- [51] — *Empirical Processes with Applications to Statistics*, Wiley, New York 1986.
- [52] N. V. Smirnov, *Probability Theory and Mathematical Statistics. Collection of Papers* (in Russian), Nauka, Moscow 1970.
- [53] W. Stute, *The oscillation behavior of empirical processes*, Ann. Probab. 10 (1982), pp. 86–107.
- [54] G. S. Watson, *Goodness-of-fit tests on a circle*, Biometrika 48 (1961), pp. 109–114.
- [55] V. V. Yurinskii (V. V. Jurinskii), *Bounds for characteristic functions of certain degenerate multidimensional distributions*, Theory Probab. Appl. 12 (1972), pp. 101–113.
- [56] R. Zitikis, *Asymptotic expansions in the Local Limit Theorem for the Generalized Cramér–von Mises–Smirnov Statistics*, Ph. D. Thesis, Vilnius University, Vilnius 1988.
- [57] — *Smoothness of distribution function of \mathcal{FL} -statistic. I*, Lithuanian Math. J. 30 (1990), pp. 231–240.
- [58] W. R. van Zwet, *Asymptotic expansions for the distribution functions of linear combinations of order statistics*, in: *Statistical Decision Theory and Related Topics. II*, S. S. Gupta and D. S. Moore (Eds.), Academic Press, New York 1977, pp. 421–436.

Fakultät für Mathematik
Universität Bielefeld
Postfach 8640
4800 Bielefeld 1, Germany

Institute of Mathematics
and Informatics
Akademijos 4
Vilnius 2600, Lithuania

Received on 20.12.1993

